

## Growth and Body Composition of Laying Hens under Different Feeding Programs up to 72 Weeks

Andréa Luciana dos Santos<sup>1\*</sup>, Douglas Emygdio de Faria<sup>2</sup>, Roselaine Ponso de Oliveira<sup>2</sup>, Mariana Pavesi<sup>2</sup>, Mayra Fernanda Rizzo Silva<sup>2</sup>, Fábio Luiz Buranelo Toral<sup>3</sup>, César Gonçalves de Lima<sup>4</sup>, Alessandro Borges Amorim<sup>1</sup>, Mayra Anton Dib Saleh<sup>5</sup>, Antonio Callejo Ramos<sup>6</sup> and Carlos Buxade Carbo<sup>6</sup>

<sup>1</sup>Institute of Agricultural Sciences and Technologies, Federal University of Mato Grosso, Rondonópolis, Mato Grosso, Brazil

<sup>2</sup>Department of Animal Science, São Paulo University, São Paulo, Brazil

<sup>3</sup>Department of Animal Science, Federal University of Minas Gerais, Minas Gerais, Brazil

<sup>4</sup>Department of Basic Sciences, São Paulo University, São Paulo, Brazil

<sup>5</sup>Department of Animal Production, Faculty of Animal Science and Veterinary Medicine, UNESP – São Paulo State University, Lageado Experimental Farm, Botucatu Campus, São Paulo, Brazil

<sup>6</sup>Department of Agrarian Production, University Politecnica of Madrid, Madrid, Spain

\*Corresponding author: Andréa Luciana dos Santos, Institute of Agricultural Sciences and Technologies, Federal University of Mato Grosso, Highway MT-270, Rondonópolis, Mato Grosso, Brazil, E-mail: [andrealfia@gmail.com](mailto:andrealfia@gmail.com)

Received date: 25 Aug 2017; Accepted date: 10 Oct 2017; Published date: 16 Oct 2017.

Citation: dos Santos AL, de Faria DE, de Oliveira RP, Pavesi M, Silva MFR, et al. (2017) Growth and Body Composition of Laying Hens under Different Feeding Programs up to 72 Weeks. J Anim Sci Res 1(1): doi <http://dx.doi.org/10.16966/2576-6457.103>

Copyright: © 2017 dos Santos AL, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

This study aimed at comparing growth and changes in body composition of laying hens under different feeding programs from hatch to 72 week old. Hens from  $W_{36}$  or Brown lineages were fed 95% ( $P_1$ ), 100% ( $P_2$ ) or 105% ( $P_3$ ) of their nutritional requirements. Six pens (replicates) with 46 hens each were used for each combination of lineage and feeding program. Growth curves were fitted using Gompertz's model. The highest body growth rate was observed for Brown-hens fed  $P_2$  at 8 week old, when the maximum growth rate and maturity rate were achieved. The same lineage of hens fed  $P_3$  presented the greatest estimated body weight at maturity.  $W_{36}$  hens had higher deposition of fat and protein throughout the 6<sup>th</sup> week at maximum growth rate compared to Brown hens. This resulted in  $W_{36}$  hens that reached body weights at maturity faster than Brown hens. Fat deposition occurred at a later age than protein deposition in both lineages and all feeding programs. The present data support the usage of 105% of the declared nutritional requirement for the best growth performance.

**Keywords:** Body Contents; Nutritional Requirements; Performance; Poultry

### Introduction

Growth of laying hens through 18-week-old is usually divided into three phases of six weeks each. Each phase is mainly characterized by determining physiological aspects: bone and muscle formation (0 to 6 weeks), feathering (6 to 12 weeks), and reproductive tract formation (12 to 18 weeks), respectively [1].

Rapid changes in poultry efficiency since the 1950's were mostly due to improved genetics, e.g. broilers [2], achieved through selection based on quantitative-genetic procedures. Nutritional requirements of birds need to be constantly re-evaluated and updated in order to design biologically meaningful and economically efficient feeding programs. Thus, accurate description of growth, meat and fat deposition patterns are pursued to achieve the above goals.

Modern poultry industry requires the establishment of precise nutritional programs in order to attain the full genetic potential of birds. Efficient deposition of protein, fat, and other nutrients in proper amounts are aimed at growth stage of birds and, especial attention must be paid to avoid excessive deposition of fat above the level that is physiologically required [3] because, fat is a nutritionally expensive tissue.

Martin et al.[4] reported that feeding programs and monitoring of growth are less studied in pullets and hens than in broilers. Prediction of body composition at different stages of growth is achieved by using growth curve models, useful for tracking changes in body composition over time, both in its chemical and physical aspects. In particular, the

model described by Gompertz et al.[5] has been valuable to describe several traits of growing birds [6-8]. Considering the aforementioned, this study aimed to compare the growth and body composition of two lineages of commercial laying hens under different nutritional programs by employing Gompertz's model.

### Materials and Methods

All experimental procedures were previously approved by the Animal Ethics Committee of this university and, in accordance with directive 2010/63/EU. Protocols for bird care and utilization, including the exit method were in compliance with regulations set forth by the host institution and funding agencies and were strictly followed throughout the trial. Birds from two genetic lineages of high-yield laying hens (Hy Line  $W_{36}$  and Hy-Line Brown) were used in the trial. Concerning the adult body weight, the hens from the  $W_{36}$  lineage are considered light and those from the Brown lineage are considered semi-heavy. Each lineage contributed with 828 birds that were raised in metal cages from 1-day-old through 72 weeks.

Chicks were individually weighed at 1-day-old and, 36 groups were formed with comparable initial average body weight, distributed among nutritional programs aiming at having similar average weights at the beginning of the trial. Each experimental unit consisted of 46 chicks. A 3×2 factorial arrangement was established in a completely randomized design. The factors were genetic lineage and nutritional program. The latter was based on the nutritional requirements set forth in breeding company commercial guides [9,10] and consisted of  $P_1$  (95% of the

nutritional requirement);  $P_2$  (100% of the nutritional requirement) and,  $P_3$  (105% of the nutritional requirement). Each combination of lineage and nutritional program had six replicates.

Animals had *ad libitum* access to feed and water. Growers and pullets were vaccinated against bronchitis, infectious bursa disease, Newcastle disease, infectious coryza and, avian pox. When the pullets were moved to the laying facilities, they were given an Intermult 6' vaccine for multiple ailments; feed and water were also available at all times. Air temperature and humidity were recorded twice daily and remained between 24.0°C-29.8°C and 56%-84% (up to 6 weeks), 16.7°C-27.8°C and 37.7%-76.6% (through 17 weeks), and 19.1°C-28.3°C and 44.7%-79.5% (through 72 weeks), respectively.

The maximum age at which birds reached sexual maturity among all treatments was 18 weeks of age. The lighting program was established according to the breeding company recommendations [9-12] for the completion of the standard nutritional program,  $P_2$ : 100% of requirements (Table 1), and respectively the requirements of HyLine manual showing that the study is within the recommended requirement [13,14]. The other programs had their requirements changed in 5% lower ( $P_1$ : 95% of requirements) and 5% above ( $P_3$ : 105% of requirements) in relation to nutritional requirements  $P_2$  (100%). A detailed description of housing facilities and husbandry procedures were reported by Santos et al. [15].

Individual body weights were taken weekly until to 6-weeks-old and then biweekly throughout the trial. Average body weights were obtained for each experimental unit and growth curves were fitted to each combination of genetic line and nutritional program, using the model proposed by Gompertz (1925):  $W_t = W_m \exp(-\exp(-b(t-t^*)))$ , where:  $W_t$  is the body weight of the bird at age  $t$ , expressed as a function of  $W_m$ ,  $W_m$  is the bird's body weight at maturity,  $b$  is the maturity constant (or maturity rate), and  $t^*$  is the age at which growth rate is at its maximum.

The chemical composition of the carcasses [16], water content [16], fat content as ethereal extract [16] (920.39), crude protein [16] (988.05), and ashes [16] (942.05) obtained based on the average data of two birds from each experimental unit (Figure 1).

At 1-day-old and on weeks 4, 6, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, and 72, two birds with body weights closest to the average of the experimental unit were euthanized. After a 24h fasting (access to water was maintained) the birds were weighed, killed by cervical dislocation and placed in an autoclave and fully mixed in an industrial type blender. An aliquot from each pair of mixed birds was dried in a convection oven at 65°C for 72h and, subjected to grinding again. Further chemical analyses were carried out to determine the ethereal extract, crude protein, and ashes contents in each sample.

Allometric relationships were calculated using the exponential function of Brody (1945) [17]:  $Y=aX^b$ , where:  $Y$ =content of the body component in grams [relations to live body weight (grams)],  $X$ =composition of carcasses in grams,  $a$ =extrapolation of  $Y$  for  $X=1$ ;  $b$ =allometric coefficient, the ratio of percentage change in  $Y$  to the corresponding percentage change in  $X$ . The slope (allometric coefficient) was calculated for each treatment versus lineage (six values), and for each carcass component (crude protein, water, ethereal extract and ashes in grams, the same unit as the unit of body weight). Thus, 24 allometric coefficients were obtained. A two-way between groups ANOVA analysis of variance by SAS [18] software was used to evaluate nutritional programs and hen's lineage interaction effects for dependent variables. Tukey's multiple range tests were applied to test significant differences between treatments ( $\alpha=0.05$ ).

## Results

Estimates of the growth curve parameters obtained using Gompertz's model is presented in table 2. The birds from the  $W_{36}$  lineage had faster



**Figure 1:** Sequence of sample preparation.

a) autoclave opening; b) two birds per experimental unit in each division; c) each division with lids; d,e) opening after autoclaving; f) birds autoclaving; g) carcasses blending; h,i) sampling carcass; j) distribution of aluminum trays inside convection oven; k) sample weighting+aluminum tray; l,m) drying the sample in a convection air oven at 65°C; n) dry sample after 72-hour; o) grinding of sample in a ball milling machine

maturity rates (parameter  $b$ ) than those from the Brown lineage, but lower values of maximum growth rate. The age at which the maximum growth rate obtained was lower in  $W_{36}$  lineage (4.5 days).

As a result, the  $W_{36}$  birds attained its mature body weight faster than those of the Brown-lineage (Figure 2). Birds from the  $W_{36}$  lineage under nutritional program  $P_3$  and  $P_2$  had higher maturity rate compared to  $P_1$ , as well as the highest body weight. Their predicted body weights at maturity were closest to the figure of 1.53 and 1.56 kg presented in the commercial management guide, respectively [10,13,14] and, the same real value was obtained at this trial, according to table 3.

Brown-lineage birds fed  $P_3$  had the highest body weight, but there was no statistical difference in maturity rate for birds that received different requirements. The latest group was the only one to achieve or exceed the target mature body weight of 1.92 kg established by the commercial management guide [9]; although, all nutritional requirements have not reached the mature body weight of 1.97 kg established by the commercial management guide[13,14].

In addition, Brown lineage birds had higher deposited protein weight than the  $W_{36}$  lineage at 8<sup>th</sup> up to 12<sup>th</sup> week, period of age at maximum growth rate, by an average of about 55.2 and 46.8 g, respectively (Table 3), across nutritional guides. Brown birds had similar protein deposition than  $W_{36}$ . Both lineages attained this maximum weight rate at essentially the same age. The nutritional programs produced very similar patterns of protein deposition within genetic lineage.

The deposited fat weight was higher for the  $W_{36}$  lineage than for the Brown lineage from day one to 16<sup>th</sup> week, similar to the pattern of protein deposition in brood and grows out. After the 60<sup>th</sup> week, the hens from the Brown-lineage had higher fat deposition than hens from  $W_{36}$  lineage. There was no interaction between the effects of lineage and feeding regimen. The

**Table 1:** Composition and calculated nutrient levels of basal feed

Ingredient (%)	Age of hens (weeks)													
	1-6		7-9		10-17		18-32		33-44		45-58		59-72	
	W <sub>36</sub>	Brown	W <sub>36</sub>	Brown	W <sub>36</sub>	Brown	W <sub>36</sub>	Brown	W <sub>36</sub>	Brown	W <sub>36</sub>	Brown	W <sub>36</sub>	Brown
Corn grains	64.618	64.754	60.801	55.697	75.202	77.386	62.877	57.796	63.686	62.332	64.179	59.871	67.072	69.181
Soybean meal (48%)	30.458	30.543	19.775	22.523	16.682	11.764	18.743	23.683	17.924	20.383	19.790	23.016	19.912	18.566
Limestone	0.937	1.112	1.076	1.139	1.166	0.974	7.876	8.970	9.709	9.300	9.710	10.550	10.248	10.262
Gluten meal (60%)	-	-	5.350	7.209	2.303	1.301	5.000	3.000	1.469	-	1.852	1.598	-	-
Wheat bran	-	-	-	-	-	3.620	-	-	-	-	-	-	-	-
Meat and bone meal	-	-	-	-	-	3.774	-	-	-	5.270	-	-	-	-
Dicalcium phosphate	2.110	1.839	-	-	1.916	0.541	2.195	1.859	1.790	-	1.789	1.437	1.528	1.200
Salt	0.352	0.352	2.028	1.898	0.343	0.305	0.372	0.368	0.348	0.279	0.348	0.368	0.372	0.353
Soybean oil	0.928	0.854	0.347	0.370	-	-	2.468	4.002	2.000	2.082	2.000	2.843	0.455	0.035
L-Lys.HCl (78%)	0.088	0.021	3.075	3.526	0.137	-	0.116	0.045	0.052	0.043	0.054	0.028	0.015	0.050
DL-Met (99%)	0.099	0.098	0.184	0.259	0.060	-	0.081	0.069	0.143	0.094	0.028	0.039	0.123	0.103
L-Trp (98%)	-	-	-	-	-	-	0.003	0.003	-	0.017	-	-	-	-
L-Thr (98%)	0.010	0.028	0.064	0.079	-	0.035	0.065	0.005	0.096	-	-	-	0.025	-
Mineral premix	0.400 <sup>1</sup>	0.400 <sup>1</sup>	0.300 <sup>1</sup>	0.300 <sup>1</sup>	0.300 <sup>1</sup>	0.300 <sup>1</sup>	0.200 <sup>2</sup>	0.200 <sup>2</sup>	0.200 <sup>2</sup>	0.200 <sup>2</sup>	0.200 <sup>2</sup>	0.200 <sup>2</sup>	0.200 <sup>2</sup>	0.200 <sup>2</sup>
Choline chloride	-	-	-	-	-	-	-	-	-	-	0.050	0.050	0.050	0.050
Kaolin	-	-	7.000	7.000	1.891	-	-	-	0.583	-	-	-	-	-
<b>Composition (%)</b>														
ME MJ/kg	12.35	12.35	12.43	12.43	12.56	12.56	12.35	12.35	11.93	11.93	11.93	11.93	11.72	11.72
Crude protein	20.00	20.00	18.00	17.50	16.00	15.50	17.50	18.00	15.50	17.50	15.25	17.00	15.00	14.54
Calcium	1.00	1.00	1.00	1.00	1.00	1.00	3.65	4.00	4.10	4.25	4.25	4.50	4.40	4.32
Digestible P	0.50	0.45	0.47	0.43	0.45	0.42	0.50	0.44	0.46	0.40	0.42	0.36	0.38	0.35
Sodium	0.18	0.18	0.17	0.18	0.17	0.18	0.18	0.18	0.17	0.18	0.17	0.18	0.17	0.16
Methionine	0.48	0.48	0.43	0.41	0.39	0.32	0.48	0.46	0.40	0.46	0.38	0.41	0.37	0.38
Methionine+Cystine	0.75	0.75	0.70	0.71	0.66	0.58	0.82	0.76	0.70	0.76	0.56	0.61	0.65	0.62
Lysine	1.15	1.10	0.96	0.90	0.85	0.66	0.88	0.93	0.82	0.93	0.78	0.89	0.76	0.75
Threonine	0.73	0.73	0.67	0.55	0.61	0.52	0.68	0.65	0.66	0.65	0.54	0.61	0.62	0.58
Tryptophan	0.20	0.20	0.18	0.19	0.16	0.18	0.18	0.20	0.17	0.20	0.16	0.19	0.17	0.16
Linoleic acid	1.95	1.92	3.04	3.24	1.59	1.59	2.75	3.48	2.48	2.48	2.48	2.87	1.642	1.448

Composition and calculated nutrient levels of basal feed established according to recommendations from breeding company (HyLine International, 2005 – 2007 a,b) for the completion of the standard nutritional program, P<sub>2</sub>: 100% used for preparation of experimental diets for hens in the study (%) [1]. Supplementation of vitamins, minerals and additives per kg of product: Vitamin A=1,900,000 I.U.; Vitamin D<sub>3</sub>=400,000 I.U., Vitamin E=3,325 mg, Vitamin K<sub>3</sub>=0.70 mg, Vitamin B<sub>2</sub>=1,000 mg, Vitamin B<sub>12</sub>=2,671 µg; Calcium pantothenate=2,850 mg; Methionine=200 mg, Niacin=6,676 mg, Choline chloride=76,000 mg, Selenium=40 mg, Copper=2,000 mg, Iron=16,800 mg, Zinc=16,226 mg, Manganese=14,300 mg, Iodine=150 mg, Antioxidant (B.H.T.: butylated hydroxy toluene)=600 mg, Excipient q.s.=1,000 g [2]. Supplementation of vitamins, minerals and additives per kg of product: Vitamin A=1,800,000 I.U.; Vitamin D<sub>3</sub>=370,000 I.U., Vitamin E=150 mg, Vitamin K<sub>3</sub>=10 mg, Vitamin B<sub>2</sub>=80 mg, Vitamin B<sub>12</sub>=110 µg; Calcium pantothenate=80 mg; Methionine=13 g; Niacin=200 mg, Choline chloride=2,500 mg, Se=3 mg; Cu=120 mg, Fe=1,000 mg, Zn=1,475 mg, Mn=1,236 mg, I=13 mg, Antioxidant (B.H.T.: butylated hydroxytoluene) =600 mg; Growth additive=1,000 g; F (maximum)=500 mg; Excipient q.s.=1,000 g.

**Table 2:** Parameter estimates of growth curve obtained using Gompertz's equation for W<sub>36</sub> and Brown laying hens submitted to different nutritional programs.

Equation Parameters	Lineages and nutritional programs					
	W <sub>36</sub>			Brown		
	P <sub>1</sub>	P <sub>2</sub>	P <sub>3</sub>	P <sub>1</sub>	P <sub>2</sub>	P <sub>3</sub>
	Trait: Live body weight					
W <sub>m</sub> (g) <sup>a</sup>	1,450.8 <sup>Bb</sup>	1,459.6 <sup>Bab</sup>	1,515.7 <sup>Ba</sup>	1,862.5 <sup>Ac</sup>	1,890.5 <sup>Ab</sup>	1,948.0 <sup>Aa</sup>
b (g/wk) <sup>b</sup>	0.1832 <sup>Aa</sup>	0.1685 <sup>Ab</sup>	0.1692 <sup>Ab</sup>	0.1604 <sup>Ba</sup>	0.1650 <sup>Aa</sup>	0.1582 <sup>Ba</sup>
t(wk) <sup>c</sup>	7.5374 <sup>Ba</sup>	7.8269 <sup>Ba</sup>	7.6763 <sup>Ba</sup>	8.3258 <sup>Aa</sup>	8.2737 <sup>Aa</sup>	8.3691 <sup>Aa</sup>
MG (g/wk) <sup>d</sup>	94.6 <sup>Ba</sup>	90.44 <sup>Bb</sup>	94.21 <sup>Ba</sup>	109.75 <sup>Ab</sup>	114.63 <sup>Aa</sup>	113.17 <sup>Aa</sup>

P<sub>1</sub>: 95%, P<sub>2</sub>: 100%, and P<sub>3</sub>: 105% of nutritional requirements; <sup>a</sup>W<sub>m</sub>: weight at maturity; <sup>b</sup>b: maturity constant (maturity rate); <sup>c</sup>t: age at maximum growth rate; <sup>d</sup>MG: maximum growth rate [it is obtained from the derivative (dW<sub>m</sub>/dt)]. Means followed by different capital letters on lines between different HyLine lineages, within each nutritional requirements differ statistically (P<0.05). Means followed by different lowercase in lines between different nutritional requirements within each HyLine lineages differ statistically (P<0.05).

weight at maturity, as well as the maximum growth rate, was higher for Brown-birds than  $W_{36}$  birds [Figure 2(a), (b), (c)], that however, reached it earlier than in the Brown-birds. In both lineages, a trend was observed for the age at deposited fat content that did not increase the birds' body weight receiving a higher fraction than recommended nutrition intake (Table 3, Table 4).

Brown birds had similar ( $P>0.05$ ) weight ash content than  $W_{36}$ . In each lineage, the content of ashes was also similar despite the increased

nutritional intake and this increase was steeper in the Brown birds. Brown birds had similar ( $P>0.05$ ) ash content than  $W_{36}$ , despite of Brown birds had a steeper intake increase. Water deposition was higher for birds from  $W_{36}$  lineage in the initial growth phase than around 16 weeks of age ( $P<0.05$ ). There was no difference ( $P>0.05$ ) in age among nutritional programs and also, there was a trend for increased water content up to 6<sup>th</sup> week of age in birds receiving the higher percentage of recommended nutritional requirement (Table 3).

**Table 3:** Live body weight and chemical composition of  $W_{36}$  and Brown laying hens submitted to different nutritional programs.

Nutrient content (g/kg)	Nutritional Programs (NP)			Lineages (L)		CV (%)
	$P_1$	$P_2$	$P_3$	$W_{36}$	Brown	
<b>Day 1</b>						
Body weight	37.54	37.29	37.28	37.64 <sup>a</sup>	37.10 <sup>b</sup>	1.05
Crude protein	293.29	342.55	324.94	317.14	323.38	12.59
Crude fat	87.40	98.94	87.78	97.06 <sup>a</sup>	85.68 <sup>b</sup>	11.07
Ash	30.10	31.72	30.57	31.56	30.03	8.56
Watercontent	589.21	526.79	556.71	554.23	560.91	8.24
<b>6<sup>th</sup> week</b>						
Body weight	417.36 <sup>b</sup>	417.52 <sup>b</sup>	432.40 <sup>a</sup>	392.37 <sup>b</sup>	452.53 <sup>a</sup>	2.43
Crude protein	250.68	216.45	229.69	248.46 <sup>a</sup>	216.09 <sup>b</sup>	10.22
Crude fat	109.32 <sup>a</sup>	84.65 <sup>b</sup>	98.25 <sup>ab</sup>	115.58 <sup>a</sup>	79.23 <sup>b</sup>	9.55
Ash	33.24	31.80	35.77	33.05	34.16	8.96
Watercontent	606.76 <sup>b</sup>	667.10 <sup>a</sup>	636.29 <sup>ab</sup>	602.91	670.52	4.78
<b>16<sup>th</sup> week</b>						
Body weight	1292.55 <sup>b</sup>	1308.80 <sup>ab</sup>	1331.11 <sup>a</sup>	1187.61 <sup>b</sup>	1434.04 <sup>b</sup>	1.99
Crude protein	159.97	201.50	228.88	223.70 <sup>a</sup>	169.87 <sup>b</sup>	11.01
Crude fat	139.79 <sup>a</sup>	138.38 <sup>ab</sup>	119.25 <sup>b</sup>	136.66	128.29	11.73
Ash	33.43	32.49	31.99	32.89	32.38	8.14
Watercontent	687.35 <sup>a</sup>	627.63 <sup>b</sup>	599.34 <sup>b</sup>	606.75 <sup>b</sup>	669.46 <sup>a</sup>	4.19
<b>32<sup>th</sup> week</b>						
Body weight	1522.16 <sup>b</sup>	1568.18 <sup>b</sup>	1630.65 <sup>a</sup>	1366.57 <sup>b</sup>	1780.75 <sup>a</sup>	3.87
Crude protein	164.51	174.71	190.33	172.24	180.80	12.53
Crude fat	155.77	172.82	160.26	153.52	172.38	17.33
Ash	39.39	35.63	39.61	37.37	39.05	11.50
Watercontent	640.32	616.85	609.80	636.87	607.77	9.02
<b>72<sup>th</sup> week</b>						
Body weight	1727.93 <sup>b</sup>	1744.87 <sup>b</sup>	1795.11 <sup>a</sup>	1569.06 <sup>b</sup>	1942.88 <sup>a</sup>	2.42
Crude protein	177.65	206.12	190.93	188.84	194.30	10.15
Crude fat	136.67 <sup>b</sup>	181.25 <sup>a</sup>	170.35 <sup>ab</sup>	161.75 <sup>b</sup>	163.77 <sup>a</sup>	16.06
Ash	43.57	49.04	40.08	42.51	45.95	14.44
Watercontent	592.11	563.59	598.64	606.90	562.66	14.64
Interaction NP×L	NS	NS	NS	NS	NS	NS


$P_1$ : 95%,  $P_2$ : 100% and  $P_3$ : 105% of nutritional requirements; CV=coefficient of variation. Means followed by different lowercase in lines differ statistically ( $P<0.05$ ). NS=interaction not significant ( $P>0.05$ ).

**Table 4:** Coefficients for allometric relations between chemical compositions of carcasses and live body weight of  $W_{36}$  and Brown laying hens submitted to different nutritional programs.

	$W_{36}$								
	$P_1$			$P_2$			$P_3$		
	b	± SE	R <sup>2</sup>	b	± SE	R <sup>2</sup>	b	± SE	R <sup>2</sup>
<b>Water</b>	0.994	0.020	0.993	1.022	0.026	0.989	0.975	0.021	0.993
<b>Protein</b>	0.933	0.031	0.982	0.934	0.031	0.982	0.934	0.031	0.983
<b>Fat</b>	1.189	0.055	0.966	1.177	0.062	0.957	1.246	0.045	0.980
<b>Ash</b>	1.039	0.022	0.993	1.038	0.022	0.988	1.038	0.021	0.993
<b>Brown</b>									
	$P_1$			$P_2$			$P_3$		
	b	± SE	R <sup>2</sup>	b	± SE	R <sup>2</sup>	b	± SE	R <sup>2</sup>
<b>Water</b>	0.996	0.017	0.995	0.984	0.019	0.994	1.009	0.0178	0.995
<b>Protein</b>	0.941	0.027	0.987	0.942	0.027	0.987	0.941	0.026	0.987
<b>Fat</b>	1.230	0.053	0.969	1.259	0.045	0.979	1.210	0.052	0.971
<b>Ash</b>	1.092	0.026	0.991	1.091	0.026	0.991	1.091	0.025	0.991

\*\*\*significance of coefficients for allometric relations between chemical composition of carcasses and live body weight in grams ( $p<0.001$ ).  $P_1$ : 95%,  $P_2$ : 100% and  $P_3$ : 105% of nutritional requirements; b: maturity constant (maturity rate); SE: standard error of the mean; R<sup>2</sup>: coefficient of determination





**Figure 2:** Growth rate curves throughout 72 weeks for W36 (♀) and Brown (♀) laying hens receiving 95% (a), 100% (b) or 105% (c) of recommended nutritional requirements.

## Discussion

The laying hens' growth management before puberty is essential to a profitable egg production, because it prepares the future laying hens for efficient reproductive output. The study of growth patterns of components of live weight allows a deeper understanding of how nutrients are being used and stored by the birds; imbalances could affect several components of total egg output like delaying the onset of lay, a lower peak production or a lower persistency of lay.

In the conditions of this trial, nutritional program P<sub>1</sub> provided inadequate nutrition to achieve the target body weights specified by the breeder company. Increasing the amount of feed intake by 5% proved a suitable adjustment to achieve the targets set by the breeder company. The birds were raised in environmental conditions that were slightly warmer than the prescribed ones for these lineages, with an exception described ahead. This may have contributed to the finding that additional 5% of feed intake provided by nutritional program P<sub>3</sub> was necessary for growers to attain their target body weights before puberty.

The inflexion point of growth curves for body weight occurred later for the Brown laying hens (Table 3, Figure 2). In part, this is due to their larger body weight at maturity and, due to the nutritional program for growing layers does not pursue rapid growth but a steady increase in body weight towards the ideal body size for optimal egg production. Heavier birds will mature later and their growth potential must be delayed to allow adequate bone and organ formation and, for optimal egg production [19]. Maximum growth rate was attained quite early, result that corroborates previous findings by Braccini Neto et al.[20] and Neme et al.[1]. Winsor, et al.[21] reported the mathematical properties of Gompertz's model and pointed out some of its uses and limitations. The model is usually suitable to describe growth patterns when the inflexion point occurs at a relatively early stage, around 35% to 40% of mature weight, which was around 32 weeks old in this trial; at this age, the hens had reached around 90% of  $W_m$ .

A portion of the heavier predicted body weights of Brown hens was deposited as protein. Altering the nutritional program did not affect the absolute value of protein content within each lineage, suggesting that selection has led the birds to have a gap on the amount of this tissue and that any additional amino acids would not participate in further anabolic process of muscle tissue. The change in body composition reflected in additional ashes, but not in fat, water and protein.

Protein is not the main form of energy storage, because its deposition is limited by the growth potential of birds; however, fat deposition is not limited by it. This developmental limitation has also evaluative importance, since lipid reserves are the main stored tissue to be mobilized; protein will only be used as an energy source after the stored fat and glycogen are closed to depletion [22]. The amount of muscle in laying hens, selected for efficiency in egg production, is limited to what is necessary to keep body structural soundness. Our data show that neither less nor additional protein is stored in the body when nutritional intake is changed; the

excess in available amino acids is likely to be channelled to protein formation in the egg. Content deposition fat occurred at a later age than protein deposition regardless of genetic lineage and nutritional program, according to coefficients for allometric relations between body component and live weight of laying hens (Table 4). The fat content deposition presented significant difference ( $P < 0.05$ ) at 72<sup>th</sup> week, occurring later for P<sub>2</sub> and P<sub>3</sub>, which may be related to a higher availability of metabolites from fat catabolism after peak growth for all other carcass components. The fat content in chickens is partially controlled by genes located on several chromosomes [23-26].

Although most studies have been conducted with lineages of broilers, there is experimental evidence for the same type of genetic control in crosses between broilers and layers [27,28], suggesting that expression of genes that control biochemical pathways responsible for fat deposition is conserved across genetic lineages. Selection for body fat content in layers is likely to receive less attention than in broilers in selection indexes due to the unequal importance of this trait in both industries.

Any selection pressure focused on reduction of body fat will mostly likely be reflected in the form of a correlated response to selection. Raising birds on a non-standard nutritional program (e.g. P<sub>3</sub>) may produce changes in the amount of fat body content. Temperature in the facility was relatively low between weeks 44 and 72 (low average weekly readings were between 14.9°C and 19.1°C). Starting on week 44, there was a noticeable reduction in fat deposition regardless of genetic lineage or nutritional program. There was no adjustment in allowance of feed, because of temperature, so the dropping on body fat deposition rate may be partially attributed to lower environmental temperatures.

Laying hens express their full genetic potential at temperatures falling in the thermal neutral zone, between 19°C and 27°C [29]. At colder temperatures, fat is mobilized to generate body heat, and at warmer temperatures, the energy is wasted to cool the body even though this is a physiologically inefficient process once chickens lack sweat glands and, the evaporation rate is too low [30]. However, increasing the amount of fat consumed by layers in colder environmental temperatures should not be employed as a strategy, because an excess in body fat accumulation can lead to welfare problems as increased death rates, like fatty liver haemorrhagic syndrome [31].

Bone strength is a complex trait influenced by many factors [19], including nutrition. The present data support the fact that an increased feed intake resulted in larger ash content in the layers' body. The larger body size of Brown layers compared to W<sub>36</sub> layers from day one until the 16<sup>th</sup> ( $P < 0.05$ ) probably required a higher level of bone mineralization; thus, higher amounts and proportions were found as more nutrients were made available through a more abundant availability of feed.

The latter phase in deposition of ashes found for the Brown lineage is consistent with other components of live weight, as larger and less precocious birds need to accumulate more resources before finalizing

their growth in a way that supports high reproductive capacity. There may be a physiological limit to the speed of bone mass growth that requires heavier birds a longer time to attain the ideal mineralization of main bones. Selection has altered the efficiency of many physiological processes in domesticated chickens and nutrient utilization, despite a conserved digestibility process even after many generations of artificial selection, may have resulted in a changed strategy for allocation of nutrients in modern commercial chickens [32].

Layers from the two genetic lineages had distinct growth profiles and, the deposition of nutrients; also, followed different schedules according to age (Table 3, Table 4, Figure 2). There was an effect of nutritional programs on some traits (live body weight and fat content).

## Conclusion

Gompertz's model was an adequate descriptor of growth curve, because the coefficients of determination were quite high for each combination of genetic lineage and nutritional program. Rearing the birds with a nutritional program that offers 105% of the recommendation by the breeding company resulted in a better match to the hens' growth and nutrient deposition potential.

## Acknowledgment

This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) under grant number 06/01733-8.

## Disclosure statement

None of the authors has any financial or personal relationships that could inappropriately influence or bias the content of the paper.

## References

- Neme R, Sakomura NK, Fukayama EH, Freitas ER, Fialho FB, et al. (2006) Growth curves and deposition of body components in pullets of different strains. *R Bras Zootec* 35: 1091-1100.
- Havenstein GB, Ferket PR, Qureshi MA (2003) Growth, livability, and feed conversion of 1957 versus 2001 broilers when fed representative 1957 and 2001 broiler diets. *Poult Sci* 82: 1500-1508.
- Emmans GC (1996). The effect of temperature on performance of laying hens. In: Morris TR, Freeman BM (eds), *Energy requirements of poultry*, British Poultry Science, Edinburgh 79-90.
- Martin PA, Bradford GD, Gous RM (1994) A formal method of determining the dietary amino acid requirements of laying-type pullets during their growing period. *Br Poult Sci* 35: 709-724.
- Gompertz B (1825) On the nature of the function expressive of the law of human mortality, and on a new method of determining the value of life contingencies. *Philos Trans R Soc Lond B Biol Sci* 115: 513-585.
- Gous RM, Moran ET Jr, Stilborn HR, Bradford GD, Emmans GC (1999) Evaluation of the parameters needed to describe the overall growth, the chemical growth, and the growth of feathers and breast muscles of broilers. *Poult Sci* 78: 812-821.
- Hancock CE, Bradford GD, Emmans GC, Gous RM (1995) The evaluation of growth parameters of six strains of commercial broiler chickens. *Br Poult Sci* 36: 247-264.
- Hruby M, Hamre ML, Coon CN (1995) Predicting amino acid requirements for broilers at 21.1°C and 32.21°C. *J Appl Poultry Res* 4: 395-401.
- HY-LINE INTERNATIONAL 2005a. Hy-Line variety Brown: Commercial management guide 2005-2007. Hy-Line International, West Des Moines.
- HY-LINE INTERNATIONAL 2005b. Hy-Line variety W-36: Commercial management guide 2005-2007. Hy-Line International, West Des Moines.
- HY-LINE INTERNATIONAL 2007a. Hy-Line variety Brown: Commercial management guide 2007. Hy-Line International, West Des Moines.
- HY-LINE INTERNATIONAL 2007b. Hy-Line variety W-36: Commercial management guide 2007. Hy-Line International, West Des Moines.
- HY-LINE INTERNATIONAL 2013a. Hy-Line variety Brown: Commercial management guide 2013. Hy-Line International, West Des Moines.
- HY-LINE INTERNATIONAL 2013b. Hy-Line variety W-36: Commercial management guide 2013. Hy-Line International, West Des Moines.
- Santos AL (2008) Performance, growth, egg quality, body composition, bone and reproductive characteristics of hens fed different nutritional programs. Thesis (Ph.D. on Animal Sciences). São Paulo University, Pirassununga 175p.
- AOAC (1990). *Agricultural chemicals. Official methods of analysis. (15<sup>th</sup> edtn)* The Association of Official Analytical Chemists, Arlington, USA.
- Brody S. (1945) *Bioenergetics and growth. (1<sup>st</sup> edtn)*, Reinhold Publishing Corporation, New York, 1023 p.
- STATSOFT INC (1998). *Electronic statistics textbook*. StatSoft, Tulsa, USA.
- Rath NC, Huff GR, Huff WE, Balog JM (2000) Factors regulating bone maturity and strength in poultry. *Poult Sci* 79: 1024-1032.
- Braccini Neto J (1993) Study of laying hens genetic growth curves. Dissertation (MSc on Sciences). Pelotas Federal University, Brazil 1-102.
- Winsor CP (1932) The Gompertz curve as a growth curve. *Proc Natl Acad Sci USA* 18: 1-8.
- Blem CR (1990) Avian energy storage. In: Power DM (ed), *Current ornithology*, vol 7, Plenum Press, New York 59-114.
- Abasht B, Pitel F, Lagarrigue S, Le Bihan Duval E, Le Roy P, et al. (2006) Fatness QTL on chicken chromosome 5 and interaction with sex. *Genet Sel Evol* 38: 297-311.
- Ikeobi CO, Woolliams JA, Morrice DR, Law A, Windsor D, et al. (2002) Quantitative trait loci affecting fatness in the chicken. *Anim Genet* 33: 428-435.
- Jennen DG, Vereijken AL, Bovenhuis H, Crooijmans RP, Veenendaal A, et al. (2004) Detection and localization of quantitative trait loci affecting fatness in broilers. *Poult Sci* 83: 295-301.
- Jennen DG, Vereijken AL, Bovenhuis H, Crooijmans RP, Van Der Poel JJ, et al. (2005) Confirmation of quantitative trait loci affecting fatness in chickens. *Genet Sel Evol* 37: 215-228.
- Liu X, Li H, Wang S, Hu X, Gao Y, et al. (2007) Mapping quantitative trait loci affecting body weight and abdominal fat weight on chicken chromosome one. *Poult Sci* 86: 1084-1089.
- Liu X, Zhang H, Li H, Li N, Zhang Y, et al. (2008) Fine-mapping quantitative trait loci for body weight and abdominal fat traits: effects of marker density and sample size. *Poult Sci* 87: 1314-1319.
- Leeson S, Summers JD (2005). *Commercial Poultry Nutrition. 3<sup>rd</sup> edition*, University books, 398.
- Dawson WR, Whittow GC (2000). Regulation of body temperature. In: Whittow GC (ed), *Sturkie's Avian Physiology, 5<sup>th</sup> edition*, Academic Press, New York 343-390.
- Whitehead CC (2002). *Nutrition and poultry welfare. World's Poult Sci J* 58: 349-356.
- Jackson S, Diamond J (1996) Metabolic and Digestive Responses to Artificial Selection in Chickens. *Evolution* 50: 1638-1650.