

## Post-Thoracotomy Pain Management

Baturay Kansu Kazbek<sup>1</sup>, Ahmet Gokhan Gundogdu<sup>2</sup>, Tevfik Kaplan<sup>3</sup>, Bulent Kocer<sup>4</sup>, Serdar Han<sup>3</sup>, and Gultekin Gulbahar<sup>2\*</sup>

<sup>1</sup>Department of Anesthesiology and Reanimation, Ufuk University, School of Medicine, Ankara, Turkey

<sup>2</sup>Division of Thoracic Surgery, Dr. Nafiz Korez Sincan State Hospital, Ankara, Turkey

<sup>3</sup>Department of Thoracic Surgery, Ufuk University, School of Medicine, Ankara, Turkey

<sup>4</sup>Division of Thoracic Surgery, Numune Teaching and Research Hospital, Ankara, Turkey

**Corresponding author:** Gultekin Gulbahar, Division of Thoracic Surgery, Dr. Nafiz Korez Sincan State Hospital, Ankara, Turkey, Tel: 90 505 3359531; Fax: 90 312 2735151; **E-mail:** [mdgultekin@gmail.com](mailto:mdgultekin@gmail.com)

**Received date:** 02 Feb 2016; **Accepted date:** 10 Mar 2016; **Published date:** 14 Mar 2016.

**Citation:** Kazbek BK, Gundogdu AG, Kaplan T, Kocer B, Han S, et al. (2016) Post-Thoracotomy Pain Management. *J Clin Anesth Manag* 1(2): doi <http://dx.doi.org/10.16966/2470-9956.105>

**Copyright:** © 2016 Kazbek BK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

Thoracotomy is a very painful procedure which may be performed for a bunch of reasons. Pain following thoracotomy incision is multifactorial and difficult to treat. Treatment of postthoracotomy pain not only provides patient comfort but also prevents postoperative pulmonary morbidity. Chronic post-thoracotomy pain has a serious negative impact on the quality of life. Although thoracic epidural analgesia has been traditionally accepted as the gold standard, paravertebral block has been gaining popularity in recent time. This review attempts to investigate the treatment options for post-thoracotomy pain.

**Keywords:** Thoracotomy; Pain management; Analgesia; Anesthesia; Regional; Treatment efficacy

### Introduction

Thoracotomy is one of the surgical procedures most associated with severe postoperative pain. Postoperative pain can cause severe respiratory problems such as hypoxia, atelectasis and pulmonary infections. These conditions may result in dreadful situations and lead to respiratory failure [1]. Therefore, pain management is of crucial importance to minimize morbidity and mortality rates in patients who have undergone thoracotomy.

### Discussion

Pre-emptive analgesia has been proposed as an effective method of pain control. Although there are arguments over the effectiveness of this method, both systemic and regional analgesia have been used for the prevention of postthoracotomy pain. Clonidine, corticosteroids, nonsteroid antiinflammatory drugs, ketamine and pregabalin have been used as agents. Tschenko et al. [2] used pre-emptive clonidine (single dose intramuscular injection, 2 mcg/kg) and reported that it did not decrease Visual Analog Scale (VAS) scores or analgesic requirements compared to placebo. Bigler et al. [3] compared a single 25 mg/kg intravenous injection of methylprednisolone to placebo on 38 patients and reported that pre-emptive corticosteroid bolus reduced VAS scores after 4 hours and on the first day at rest and after 4 and 8 hours on the second day during cough. Application of steroids did not have a positive effect on pulmonary functions. Moreover, three patients in the corticosteroid group had to undergo reoperation. The authors stated that preoperative steroid injection is not warranted.

Ketamine, which is an N-Methyl-D-Aspartate (NMDA) receptor antagonist, has been used for pre-emptive analgesia by Ozyalçın et al. [4] in a study conducted in 2004. In this study, sixty patients were divided into 3 groups (intramuscular ketamine 1 mg/kg plus epidural saline, epidural ketamine 1 mg/kg plus intramuscular saline and finally intramuscular and epidural saline). The authors stated that postoperative epidural morphine and local anesthetic consumption was lower in the epidural ketamine group. The study does not report any side effects attributable to epidural ketamine usage.

Kinney et al. [5] compared gabapentin/pregabalin as pre-emptive analgesia for post-thoracotomy pain in 2012. The patients undergoing thoracotomy received 600 mg gabapentin 2 hours before the operation and were given continuous epidural infusion of gabapentin combined with ketorolac and acetaminophen. The authors stated that pre-emptive gabapentin did not reduce pain scores or opioid consumption following elective thoracotomy.

Although the efficacy of pre-emptive analgesia has been subject to debate, pre-emptive analgesia is believed to decrease the incidence of postoperative pain syndromes by changing the central processing of painful stimuli [6-9]. Pre-emptive epidural local anesthetic with or without opioids have also been used for the management of pain following thoracotomy. This method has been traditionally compared to systemic analgesia. Pre-emptive thoracal treatments offer the advantage of avoiding common postoperative side effects often associated with systemic analgesic treatment such as nausea, vomiting, sedation and difficulty in ambulation in the early postoperative period. However, regional blocks have their own morbidities. For example, pre-emptive single dose bolus of intrathecal opioids have been shown to increase the incidence of urinary retention despite the fact that they reduce the pain scores in the postoperative period [10]. Numerous studies, which compare different forms of regional pre-emptive analgesia to systemic analgesia, have been carried out. These studies have compared thoracic epidural, thoracic paravertebral, intercostal, and intrapleural blocks to systemic analgesia.

Pre-emptive thoracal epidural applications have been extensively reviewed. Thoracic epidural analgesia (TEA) is considered as the gold standard [11]. It has also been shown to be superior to conventional analgesia models in postthoracotomy pain [12,13]. In a study conducted by Ertürk et al. [14] co-infusion of levobupivacaine and fentanyl following initiating bolus injection was compared to that of saline. It was reported that local anesthetic and opioid combination decreased VAS scores at rest and coughing state. The authors did not report any differences between side effects. In a similar fashion, Yegin et al. [15] compared preoperative versus

postoperative initiation of TEA and reported that pre-emptive application of thoracic analgesia significantly lowers pain scores. Although possible benefits and harms of TEA are subject to debate, it is decidedly superior to intravenous PCA application. Nausea, vomiting, possible motor block, hypotension and consequent cardiovascular problems and inadequate control of pain can be numbered among disadvantages of this technique.

Although a considerable number of studies report successful decrease in pain scores by pre-emptive analgesia yet [16-18], there are other studies which state that pre-emptive analgesia is of little or no clinical value [19-22]. Effective pre-emptive analgesia requires a total afferent block which lasts for 2 or 3 days in addition to blockade of circulating pro-inflammatory cytokines (i.e. IL-1 $\beta$ ), which is difficult to achieve *in vivo* [23].

Surgical stress causes humoral mediator release. In a study by Amr et al. [24] the researchers investigated the effects of preoperative epidural local anesthetic application on respiratory and endocrine systems. Although significant improvement was observed in respiratory functions and analgesia in the intervention group; there was no statistically significant improvement in oxygenation, cortisol or glucose release levels between the two groups. This could be due to relatively low dose of bupivacaine used in the study. This seems to be caused by the desire to avoid motor blockade in the postoperative period. Although pain scores were lower, VAS scores on coughing state were higher in the intervention group in the early postoperative period. Choosing a different local anesthetic with a lower tendency to cause motor blockade could have yielded different results.

In regional anesthesia, adjuvants to local anesthetics are widely used. However, there are studies which suggest that their use in TEA does not enhance the quality of analgesia [25-27]. Blainoel et al. [25] reported that addition of methylprednisolone to continuous epidural infusion of local anesthetics does not cause any difference in postoperative pain scores or morphine requirements. On the other hand, a recent study compared the effects of magnesium sulphate and clonidine and reported that the addition of clonidine provided a longer analgesia, followed by magnesium sulphate. Clonidine also reduced the number of additional analgesics but it also increased the incidence of sedation [28]. Similarly, a study conducted by Chia et al. [29] compared the effects of pre-emptive and intraoperative epidural neostigmine. It is reported that pre-emptive continuous neostigmine decreases epidural analgesic consumption in the first six postoperative days.

Although TEA is accepted as the gold standard for the management of post-thoracotomy pain, it may be contraindicated in some patients. Additionally, it may be associated with complications such as hypotension, urinary retention, unsuccessful blocks and neurological sequelae. Paravertebral block (PVB) has emerged as an alternative to TEA. A recent study by Dango et al. [30] reports a statistically significant but yet small and questionable decrease in VAS scores when thoracic epidural anesthesia and a combination of PVB and intrathecal opioid are used. This combination is thus suggested as a viable alternative to thoracic epidural blocks. Pintaric et al. [31] compared continuous thoracic epidural analgesia to PVB and reported that the two technics provide similar analgesia, but the patients in the PVB group require less colloids and vasopressors to maintain the target oxygen delivery index. Scarci et al. [32] investigated more than 184 papers for comparison of thoracic paravertebral and epidural blocks. They stated that in PVB groups, VAS scores at rest and coughing state are significantly lower; oximetry measurements and respiratory function test scores are better. In epidural analgesia groups, adverse effects such as urinary retention, nausea, itching and hypotension are higher and technical difficulties are more frequent. Although plasma concentrations of cortisol are elevated in both groups, this elevation is less pronounced in the PVB group. Similarly, a systematic review by Davies et al. [1] evaluated randomized controlled trials concerning studies which compared PVB to

TEA in thoracic surgery. According to the results of ten studies, although there was no statistically significant difference between pain scores in the postoperative period; pulmonary and systemic complications were more frequent in the thoracic epidural group. Moreover, technical difficulties were more commonly met in the thoracic epidural group. As a result, they recommended PVB for thoracic surgery. Richardson et al. [33] also compared PVB and TEA. Better pulmonary function test results, lower pulmonary morbidity and better oxygenation levels were observed in PVB group. In TEA group, 10 patients were omitted from the study due to the failure in epidural catheter insertion. PVB is a relatively new technique compared to TEA. Although there is no certain evidence pointing to its superiority in pain control over TEA, its lack of disadvantages attributed to TEA and possibility of catheterisation under direct visualization in the intraoperative period by the surgeon can be considered as an advantage of this technique.

Catheterization for PVB can be achieved either percutaneously before the surgery or during the operation before the closure of the thoracic cage. Still, some probable risks exist for preoperative percutaneous catheterization. In a series of 367 patients, Lönquist et al. [34] sought for PVB related clinical problems. They reported hypotension, vascular puncture and pneumothorax as possible complications. Several studies emphasize the safety of placing the catheter under direct vision for the prevention of various complications [33-37]. A randomized trial by Gulbahar et al. [38] compares the effectiveness of TEA and PVB on postoperative pain control. They report that although VAS scores, pulmonary function tests and serum cortisol levels are similar between the two groups, PVB group is associated with shorter intervention time and less iatrogenic complications in comparison of epidural and paravertebral catheterisation techniques in post-thoracotomy pain management. A recent study failed to show any differences in efficacy between bolus and controlled infusion of local anesthetics via elastomeric pumps through a catheter placed under direct visualization [39].

The commonly accepted definition of postthoracotomy pain syndrome (PTPS) is pain that recurs or persists along a thoracotomy incision at least two months following the surgical procedure [40]. It shares the characteristics of neuropathic pain (such as burning, stabbing accompanied by dysesthesia). According to Perttunen et al. [41] the incidence of long-term postthoracotomy (Table 1) pain is 80% at three months, 75% at six months and 61% at one year and nearly half of the patients suffer from debilitating pain. Moreover, patients who suffer from PTPS report unsatisfactory results from various and costly therapies [42].

Proposed mechanisms for the occurrence of chronic postthoracotomy pain are diverse and include psychological factors [43], recurrence of the primary tumor [44], incision type [45] and direct damage to intercostal nerves [46]. The observation of allodynia and hyperalgesia in the areas innervated by the intercostal nerves suggests that the nerve injury caused by retraction or the incision itself is a likely culprit for chronic post-thoracotomy pain.

Sapkota et al. [47] hypothesized that intercostal nerve injury is the main reason for postthoracotomy pain. They compared pericostal sutures technics and found that suture technic with an intercostal muscle flap which is harvested at the beginning of the operation lowered postoperative pain scores without increasing total operation time. Likewise, Leandro et al. [48] compared pericostal and transcostal sutures. They found that transcostal sutures yield lower early and late pain scores. Elshiekh et al. [49] report that muscle-sparing thoracotomy confers a one-month advantage in skeletal muscle strength and range of motion over classical posterolateral thoracotomy

Pre-emptive analgesia
<ul style="list-style-type: none"> <li>• Systemic</li> <li>• Regional</li> </ul>
Thoracic epidural analgesia
<ul style="list-style-type: none"> <li>• Single shot</li> <li>• Continuous</li> </ul>
Paravertebral block
<ul style="list-style-type: none"> <li>• Single shot</li> <li>• Continuous</li> </ul>
Intercostal nerve blocks
Radiofrequency ablation of dorsal root ganglion
Nerve stimulation
<ul style="list-style-type: none"> <li>• Spinal cord</li> <li>• Peripheral nerve (TENS)</li> </ul>
Epidural injections
<ul style="list-style-type: none"> <li>• Steroid</li> <li>• Other</li> </ul>
Oral Gabapentin
<ul style="list-style-type: none"> <li>• Preoperative</li> <li>• Postoperative</li> </ul>
Interscalene brachial plexus block
Intraoperative intercostal nerve neurolysis
Multimodal analgesia

**Table 1:** Treatment options for post-thoracotomy pain.

TEA appears to be effective at reducing the incidence of PTPS however the time of initiation does not seem to be a determinant of this incidence [50]. According to a meta-analysis published by Bong et al. [51] who investigated six studies and 458 patients; although pre-emptive thoracic anesthesia provides a significant reduction in pain scores on coughing at 24 and 48 hours, there is no significant difference concerning the incidence of chronic pain at 6 months between pre-emptive TEA (39.6%) and control groups (48.6%). The effectiveness of PVB has not been fully investigated in prevention of PTPS. Intercostal nerve blocks, local anesthetic injections and cryoanalgesia do not seem to be effective. Moreover, cryoanalgesia may increase the incidence of neuropathic symptoms [52]. Ketamine has been investigated in two randomized studies and did not decrease the incidence of PTPS [53,54]. Mac et al. [55] report that pre-emptive and 48-hour acetaminophen treatment decreases ipsilateral shoulder pain incidence following thoracotomy.

Since PTPS has a neuropathic component, gabapentin may potentially treat this clinical condition. Solak et al. [56] compared gabapentin to naproxen sodium and showed gabapentin to be effective. Invasive interventions such as nerve blocks, radiofrequency of dorsal root ganglion, epidural injections, spinal cord and peripheral nerve stimulation may also be used.

In summary, effective postoperative pain control and treatment after thoracotomy, which is one of the most painful surgical procedures, is crucial in order to avoid pulmonary complications, delayed mobilization, late discharge and chronic pain. Although local anesthetics and opioids given via thoracic epidurals have long been accepted as gold standard for this purpose, paravertebral blocks are gaining popularity and offer the same level pain control with less complications.

Both thoracic surgeons and pain specialists should review the treatment options and tailor the chosen treatment modality for every patient's unique requirements while taking patient's decisions and other medical conditions into consideration.

## References

- Davies RG, Myles PS, Graham JM (2006) A comparison of the analgesic efficacy and side-effects of paravertebral vs epidural blockade for thoracotomy—a systematic review and meta-analysis of randomized trials. *Br J Anaesth* 96: 418-426.
- Tschernko EM, Klepetko H, Gruber E, Kritzing M, Klimscha W, et al. (1998) Clonidine added to the anesthetic solution enhances analgesia and improves oxygenation after intercostal nerve block for thoracotomy. *Anesth Analg* 87: 107-111.
- Bigler D, Jonsson T, Olsen J, Brenøe J, Sander-Jensen K (1996) The effect of preoperative methylprednisolone on pulmonary function and pain after lung operations. *J Thorac Cardiovasc Surg* 112: 142-145.
- Ozyalcin NS, Yucel A, Camlica H, Dereli N, Andersen OK, et al. (2004) Effect of pre-emptive ketamine on sensory changes and postoperative pain after thoracotomy: comparison of epidural and intramuscular routes. *Br J Anaesth* 93: 356-361.
- Kinney MA, Mantilla CB, Carns PE, Passe MA, Brown MJ (2012) Preoperative gabapentin for acute post-thoracotomy analgesia: a randomized, double-blinded, active placebo-controlled study. *Pain Pract* 12: 175-183
- Dahl JB, Hansen BL, Hjortsø NC, Erichsen CJ, Møiniche S, et al. (1992) Influence of timing on the effect of continuous extradural analgesia with bupivacaine and morphine after major abdominal surgery. *Br J Anaesth* 69: 4-8.
- Pryle BJ, Vanner RG, Enriquez N, Reynolds F (1993) Can pre-emptive lumbar epidural blockade reduce postoperative pain following lower abdominal surgery? *Anaesthesia* 48: 120-123.
- Kissin I (1996) Pre-emptive analgesia. Why its effect is not always obvious. *Anesthesiology* 84: 1015-1019.
- Wilder-Smith OH (2000) Preemptive anaesthesia and surgical pain. *Prog Brain Res* 129: 505-524.
- Chaney MA (1995) Side effects of intrathecal and epidural opioids. *Can J Anaesth* 42: 891-903.
- Ng A, Swanevelder J (2007) Pain relief after thoracotomy: is epidural analgesia the optimal technique? *Br J Anaesth* 98: 159-162.
- Debreceni G, Molnár Z, Szélig L, Molnár TF (2003) Continuous epidural or intercostal analgesia following thoracotomy: a prospective randomized double-blind clinical trial. *Acta Anaesthesiol Scand* 47: 1091-1095.
- Salomäki TE, Laitinen JO, Nuutinen LS (1991) A randomized double-blind comparison of epidural versus intravenous fentanyl infusion for analgesia after thoracotomy. *Anesthesiology* 75: 790-795.
- Erturk E, Aydogdu Kaya F, Kutanis D, Besir A, Akdogan A, et al. (2014) The effectiveness of preemptive thoracic epidural analgesia in thoracic surgery. *Biomed Res Int* 2014: 673-682.
- Yegin A, Erdogan A, Kayacan N, Karsli B (2003) Early postoperative pain management after thoracic surgery; pre- and postoperative versus postoperative epidural analgesia: a randomised study. *Eur J Cardiothorac Surg* 24: 420-424.
- Sentürk M, Ozcan PE, Talu GK, Kiyan E, Camci E, et al. (2009) The effects of three different analgesia techniques on long-term postthoracotomy pain. *Anesth Analg* 94: 11-15.
- Obata H, Saito S, Fujita N, Fuse Y, Ishizaki K, et al. (1999) Epidural block with mepivacaine before surgery reduces long-term post-thoracotomy pain. *Can J Anaesth* 46: 1127-1132.
- Richardson J, Sabanathan S, Mearns AJ, Evans CS, Bembridge J, et al. (1994) Efficacy of pre-emptive analgesia and continuous extrapleural intercostal nerve block on post-thoracotomy pain and pulmonary mechanics. *J Cardiovasc Surg* 35: 219-228.
- Neustein SM, Kreitzer JM, Krellenstein D, Reich DL, Rapaport E, et al. (2002) Preemptive epidural analgesia for thoracic surgery. *Mt Sinai J Med* 69: 101-104.
- Kavanagh BP, Katz J, Sandler AN, Nierenberg H, Roger S, et al. (1994) Multimodal analgesia before thoracic surgery does not reduce postoperative pain. *Br J Anaesth* 73: 184-189.

21. Ochroch EA, Gottschalk A, Augostides J, Carson KA, Kent L, et al. (2002) Long-term pain and activity during recovery from major thoracotomy using thoracic epidural analgesia. *Anesthesiology* 97: 1234-1244.
22. Doyle E, Bowler GM (1998) Pre-emptive effect of multimodal analgesia in thoracic surgery. *Br J Anaesth* 80: 147-151.
23. Samad TA, Moore KA, Sapirstein A, Billet S, Allchorne A, et al. (2001) Interleukin-1beta-mediated induction of Cox-2 in the CNS contributes to inflammatory pain hypersensitivity. *Nature* 410: 471-475.
24. Amr YM, Yousef AA, Alzefrawy AE, Messbah WI, Saber AM (2010) Effect of preincisional epidural fentanyl and bupivacaine on postthoracotomy pain and pulmonary function. *Ann Thorac Surg* 89: 381-385.
25. Blanloeil Y, Bizouarn P, Le Teurnier Y, Le Roux C, Rigal JC, et al. (2001) Postoperative analgesia by epidural methylprednisolone after posterolateral thoracotomy. *Br J Anaesth* 87: 635-638.
26. Baron CM, Kowalski SE, Greengrass R, Horan TA, Unruh HW, et al. (1996) Epinephrine decreases postoperative requirements for continuous thoracic epidural fentanyl infusions. *Anesth Analg* 82: 760-765.
27. Hasenbos MA, Gielen MJ, Bos J, Tielbeek E, Stanton-Hicks MD, et al. (1988) High thoracic epidural sufentanil for post-thoracotomy pain: influence of epinephrine as an adjuvant—a double blind study. *Anesthesiology* 69: 1017-1022.
28. Mohammad W, Mir SA, Mohammad K, Sofi K (2015) A randomized double-blind study to evaluate efficacy and safety of epidural magnesium sulfate and clonidine as adjuvants to bupivacaine for postthoracotomy pain relief. *Anesth Essays Res* 9: 15-20.
29. Chia YY, Chang TH, Liu K, Chang HC, Ko NH, et al. (2006) The efficacy of thoracic epidural neostigmine infusion after thoracotomy. *Anesth Analg* 102: 201-208.
30. Dango S, Harris S, Offner K, Hennings E, Priebe HJ, et al. (2013) Combined paravertebral and intrathecal vs thoracic epidural analgesia for post-thoracotomy pain relief. *Br J Anaesth* 110: 443-449.
31. Pintaric TS, Potocnik I, Hadzic A, Stupnik T, Pintaric M, et al. (2011) Comparison of continuous thoracic epidural with paravertebral block on perioperative analgesia and hemodynamic stability in patients having open lung surgery. *Reg Anaesth Pain Med* 36: 256-260.
32. Scarci M, Joshi A, Attia R (2010) In patients undergoing thoracic surgery is paravertebral block as effective as epidural analgesia for pain management? *Interact Cardiovasc Thorac Surg* 10: 92-96.
33. Richardson J, Sabanathan S, Jones J, Shah RD, Cheema S, et al. (1999) A prospective, randomized, comparison of preoperative and continuous balanced epidural and paravertebral bupivacaine on post-thoracotomy pain, pulmonary functions and stress responses. *Br J Anaesth* 83: 387-392.
34. Lönnqvist PA, Mackenzie J, Soni AK, Conacher ID (1995) Paravertebral blockade. Failure rate and complications. *Anaesthesia* 50: 813-815.
35. Fibla JJ, Molins L, Mier JM, Sierra A, Vidal G (2008) Comparative analysis of analgesic quality in the postoperative of thoracotomy: paravertebral block with bupivacaine 0.5% vs ropivacaine 0.2%. *Eur J Cardiothorac Surg* 33: 430-434.
36. Kaiser AM, Zollinger A, De Lorenzi D, Largiadèr F, Weder W (1998) Prospective, randomized comparison of extrapleural versus epidural analgesia for postthoracotomy pain. *Ann Thorac Surg* 66: 367-372.
37. Karmakar MK, Booker PD, Franks R, Pozzi M (1996) Continuous extrapleural paravertebral infusion of bupivacaine for post-thoracotomy analgesia in young infants. *Br J Anaesth* 76: 811-815.
38. Gulbahar G, Kocer B, Muratli SN, Yildirim E, Gulbahar O, et al. (2010) A comparison of epidural and paravertebral catheterisation techniques in post-thoracotomy pain management. *Eur J Cardiothorac Surg* 37: 467-472.
39. Fibla JJ, Molins L, Mier JM, Hernandez J, Sierra A (2015) A randomized prospective study of analgesic quality after thoracotomy: paravertebral block with bolus versus continuous infusion with an elastomeric pump. *Eur J Cardiothorac Surg* 631-635.
40. Koehler RP, Keenan RJ (2006) Management of postthoracotomy pain: acute and chronic. *Thorac Surg Clin* 16: 287-297.
41. Perttunen K, Tasmuth T, Kalso E (1999) Chronic pain after thoracic surgery: a follow-up study. *Acta Anaesthesiol Scand* 43: 563-567
42. Conacher ID (1992) Therapists and therapies for post-thoracotomy neuralgia. *Pain* 48: 409-412.
43. Bachiocco V, Morselli-Labate AM, Rusticali AG, Bragaglia R, Mastroianni M, et al. (1990) Intensity, latency and duration of post-thoracotomy pain: relationship to personality traits. *Funct Neurol* 5: 321-332.
44. Keller SM, Carp NZ, Levy MN, Rosen SM (1994) Chronic post thoracotomy pain. *J Cardiovasc Surg* 35: 161-164.
45. Khan IH, McManus KG, McCraith A, McGuigan JA (2000) Muscle sparing thoracotomy: a biomechanical analysis confirms preservation of muscle strength but no improvement in wound discomfort. *Eur J Cardiothorac Surg* 18: 656-661.
46. Rogers ML, Henderson L, Mahajan RP, Duffy JP (2002) Preliminary findings in the neurophysiological assessment of intercostal nerve injury during thoracotomy. *Eur J Cardiothorac Surg* 21: 298-301.
47. Sapkota R, Shrestha UK, Sayami P (2013) Intercostal muscle flap and intracostal suture to reduce post-thoracotomy pain. *Asian Cardiovasc Thorac Ann* 22: 706-711.
48. Leandro JD, Rodrigues OR, Slaets AF, Schmidt Jr AF, Yaekashi ML, et al. (2014) Comparison between two thoracotomy closure techniques: postoperative pain and pulmonary function. *J Bras Pneumol* 40: 389-396.
49. Elshiekh MA, Lo TT, Shipolini AR, McCormack DJ (2013) Does muscle-sparing thoracotomy as opposed to posterolateral thoracotomy result in better recovery? *Interact Cardiovasc Thorac Surg* 16: 60-67.
50. Khelemsky Y, Noto CJ (2012) Preventing post-thoracotomy pain syndrome. *Mt Sinai J Med* 79: 133-139.
51. Bong CL, Samuel M, Ng JM, Ip-Yam C (2005) Effects of preemptive epidural analgesia on post-thoracotomy pain. *J Cardiothorac Vasc Anesth* 19: 786-793.
52. Ju H, Feng Y, Yang BX, Wang J (2008) Comparison of epidural analgesia and intercostal nerve cryoanalgesia for post-thoracotomy pain control. *Eur J Pain* 12: 378-384.
53. Dualé C, Sibaud F, Guastella V, Vallet L, Gimbert YA (2009) Perioperative ketamine does not prevent chronic pain after thoracotomy. *Eur J Pain* 13: 497-505.
54. Ryu HG, Lee CJ, Kim YT, Bahk JH (2011) Preemptive low-dose epidural ketamine for preventing chronic postthoracotomy pain: a prospective double-blinded, randomized clinical trial. *Clin J Pain* 27: 304-308.
55. Mac TB, Girard F, Chouinard P, Boudreault D, Lafontaine ER, et al. (2005) Acetaminophen decreases early post-thoracotomy ipsilateral shoulder pain in patients with thoracic epidural analgesia: a double-blind placebo-controlled study. *J Cardiothorac Vasc Anesth* 19: 475-478.
56. Solak O, Metin M, Esme H, Solak O, Yaman M, et al. (2007) Effectiveness of gabapentin in the treatment of chronic post-thoracotomy pain. *Eur J Cardiothorac Surg* 32: 9-12.