

# Primer Design for the Identification of Ten Oral *Actinomyces* Species Using Multiplex PCR

Taira Kobayashi<sup>1\*</sup>, Satoshi Uchibori<sup>1</sup>, Osamu Tsuzukibashi<sup>2</sup>, Chihiro Uezato<sup>1</sup>, Haruhiko Goto<sup>1</sup>, Chiho Mashimo<sup>3</sup>, Takayuki Nambu<sup>3</sup>, Koji Umezawa<sup>4</sup> and Mitsuhiro Ohta<sup>5</sup>

<sup>1</sup>Department of Crown Bridge Prosthodontics, Nihon University, School of Dentistry at Matsudo, Chiba, Japan

<sup>2</sup>Department of Oral Microbiology, Nihon University, School of Dentistry at Matsudo, Chiba, Japan

<sup>3</sup>Department of Bacteriology, Osaka Dental University, Osaka, Japan

<sup>4</sup>Department of Special Needs Dentistry, Nihon University, School of Dentistry at Matsudo, Chiba, Japan

<sup>5</sup>Department of Oral Diagnostics, Nihon University, School of Dentistry at Matsudo, Chiba, Japan

\*Corresponding author: Taira Kobayashi, Department of Crown Bridge Prosthodontics, Nihon University, School of Dentistry at Matsudo, Chiba, Japan, Tel: +81-47-360-9383; E-mail: [kobayashi.taira@nihon-u.ac.jp](mailto:kobayashi.taira@nihon-u.ac.jp)

Received: 28 Feb, 2018 | Accepted: 14 Mar, 2018 | Published: 23 Mar, 2018

**Citation:** Kobayashi T, Uchibori S, Tsuzukibashi O, Uezato C, Goto H, et al. (2018) Primer Design for the Identification of Ten Oral *Actinomyces* Species Using Multiplex PCR. Int J Dent Oral Health 4(1): [dx.doi.org/10.16966/2378-7090.249](http://dx.doi.org/10.16966/2378-7090.249)

**Copyright:** © 2018 Kobayashi T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

**Background:** *Actinomyces* is one of the predominant genera in the oral cavity. Oral *Actinomyces* species play a central role in the initial stages of biofilm formation on teeth; however, limited information is currently available on the distribution of individual species in different sites or clinical conditions. Moreover, a suitable method has yet to be developed to identify oral *Actinomyces* species because of the phenotypic and genetic similarities between these microorganisms.

**Objective:** *Actinomyces naeslundii*, *A. odontolyticus*, *A. oris*, *A. georgiae*, *A. gerencseriae*, *A. graevenitzii*, *A. dentalis*, *A. johnsonii*, *A. israelii*, and *A. meyeri* among the genus *Actinomyces* are regarded as normal human oral *Actinomyces* species. The purpose of the present study was to design primers to identify oral *Actinomyces* species using multiplex PCR.

**Methods:** Polymerase chain reaction (PCR) primers were designed based on partial sequences of the 16S rDNA genes of the representative oral *Actinomyces* species. The 16S rDNA genes of the representative oral *Actinomyces* species were obtained from the DNA Data Bank of Japan, and a multiple sequence alignment analysis was performed with the CLUSTAL W program. Homology among the primers selected for oral *Actinomyces* species and their respective 16S rRNA sequences was confirmed by a BLAST search.

**Results:** These primers were able to distinguish each oral *Actinomyces* species and did not display cross-reactivity with representative oral bacteria or other *Actinomyces* species. Moreover, we developed a multiplex PCR method with the ability to identify and differentiate oral *Actinomyces* species (i.e., *A. naeslundii*, *A. johnsonii*, *A. oris*, *A. odontolyticus*, *A. israelii*, *A. georgiae*, *A. dentalis*, *A. graevenitzii*, *A. gerencseriae*, and *A. meyeri*) using only two PCR tubes per sample.

**Conclusion:** The present results indicate that our multiplex PCR method with these primers is useful for identifying the representative oral *Actinomyces* species. This method is easy because the use of Mighty Amp DNA Polymerase Ver.2 (Takara) means that DNA extraction may be avoided, and species identification using this method only takes approximately 2 hours. Thus, the method described herein will allow the prevalence of oral *Actinomyces* species and their involvement in oral infections to be fully clarified in future studies.

**Keywords:** Genus *Actinomyces*; Multiplex PCR; Oral cavity; 16S rDNA

## Introduction

Although the genus *Actinomyces* was already described in 1919, many new species have recently been discovered. The genus *Actinomyces* currently comprises 47 species and 2 subspecies (<http://www.bacterio.net/actinomyces.html>). *Actinomyces* consist of Gram-positive, anaerobic, and aerotolerant, non-spore-forming, non-motile pleomorphic rods with various degrees of branching. *Actinomyces* species are frequently found as members of the normal micro-flora, particularly in the oral cavity of humans; however, they are also etiological agents in infections, such as in classical actinomycosis, human bite wounds and abscesses at different body sites, eye infections, and oral, genital, and urinary tract infections [1,2]. The detection of these microorganisms in clinical specimens is important because it may affect the prognosis and management of patients; however, difficulties are associated with identification by conventional biochemical methods.

Ten species (i.e., *Actinomyces naeslundii*, *A. odontolyticus*, *A. oris*, *A. georgiae*, *A. gerencseriae*, *A. graevenitzii*, *A. dentalis*, *A. johnsonii*, *A. israelii*, and *A. meyeri*) among the genus *Actinomyces* are regarded as normal human oral bacteria [3]. Their distribution in different oral sites and their role in common oral afflictions, particularly periodontal diseases, remains controversial. Previous studies successfully induced destructive periodontitis in gnotobiotic rodents with *A. naeslundii* [4], suggesting a potential role for these organisms in periodontal diseases. However, an elevated level of these microorganisms in inactive lesions has also been reported [5]. Recent studies revealed that *Actinomyces* species are highly prevalent in supra- and sub-gingival plaques in adult periodontitis [6] and gingivitis [7]. Therefore, the data available on the role of *Actinomyces* species in periodontal diseases are equivocal. In addition to periodontal diseases, these microorganisms have been implicated in the pathogenesis of root surface caries and refractory periapical actinomycosis [8]. Therefore, the precise detection and speciation of these microorganisms is a prerequisite for the clarification of their pathogenic potential and possibly therapeutic interventions for the aforementioned disease entities.

The accurate identification and enumeration of *Actinomyces* species are required in order to clarify their role in oral ecology and dental diseases. Although conventional biochemical assays are used to identify *Actinomyces* species, they are often imprecise due to the phenotypic variations displayed by these bacteria. Although a sequence analysis of several target genes is the most reliable method, it is expensive, laborious, and time-consuming. Thus, a simple and more reliable assay for identifying oral *Actinomyces* species is desired. The purpose of the present study was to design primers for the identification of oral *Actinomyces* species using multiplex PCR.

## Materials and Methods

### Bacterial strains and Culture conditions

The following bacterial strains were used in the present study: *A. naeslundii* ATCC 12104, *A. oris* ATCC 27044, *A. johnsonii* JCM 16129, *A. odontolyticus* ATCC 17929, *A. israelii* ATCC 12102, *A. georgiae* DSM 6843, *A. dentalis* DSM 19115, *A. graevenitzii* DSM 15540, *A. gerencseriae* JCM 12963, *A. meyeri* ATCC 35568, *A. viscosus* ATCC 15987, *Streptococcus oralis* ATCC 10557, *S. salivarius* JCM 5707, *S. anginosus* ATCC 11391, *S. mutans* NCTC 10449, *S. sobrinus* ATCC 33478, *Rothia dentocariosa* JCM 3067, *Rothia mucilaginosa* JCM 10910, *Corynebacterium matruchotii* ATCC 14266, *C. durum* ATCC 33449, *Neisseria sicca* ATCC 29256, *Aggregatibacter actinomycetemcomitans* ATCC 33384, *Staphylococcus aureus* JCM 2874, and *S. epidermidis* ATCC 2414. These strains were maintained by cultivating them in brain heart infusion agar (BHI; Difco Laboratories, Detroit, Mich.) supplemented with 1% yeast extract (BHIY). Bacteria were cultured in BHIY broth at 37 °C for 24 h under anaerobic conditions (85% N<sub>2</sub>, 10% H<sub>2</sub>, 5% CO<sub>2</sub>) for members of the genus *Actinomyces* and in a 5% CO<sub>2</sub> atmosphere for other representative oral bacteria.

### Design of species-specific primers for ten oral *Actinomyces* species

The 16S rRNA sequences of *A. naeslundii* (accession no. AB618790), *A. johnsonii* (AB545933), *A. oris* (AB545935), *A. odontolyticus* (AB818950), *A. israelii* (AB849123), *A. georgiae* (X80413), *A. dentalis* (AJ697609), *A. graevenitzii* (AJ540309), *A. gerencseriae* (X80414), and *A. meyeri* (X82451) were obtained from the DNA Data Bank of Japan (DDBJ; Mishima, Japan), and a multiple sequence alignment analysis was performed with the CLUSTAL W program; i.e., the 16S rRNA sequences of ten oral *Actinomyces* species were aligned and analyzed. Homology among the primers selected for oral *Actinomyces* species and their respective 16S rRNA sequences was confirmed by a BLAST search.

### Development of a multiplex PCR method using designed primers

In the present study, ten oral *Actinomyces* species were divided into two groups for a multiplex PCR method. Group A consisted of *A. naeslundii*, *A. johnsonii*, *A. oris*, *A. odontolyticus*, and *A. israelii*, and group B consisted of *A. georgiae*, *A. dentalis*, *A. graevenitzii*, *A. gerencseriae*, and *A. meyeri*. Bacterial cells were cultured in BHIY broth for 24 h, and 1-ml samples were then collected in micro centrifuge tubes and resuspended at a density of 1.0 McFarland standard (approximately 10<sup>7</sup> colony-forming units (CFU)/ml) in 1 ml of sterile distilled water. A total of 3.6 µl of the suspension was then used as a PCR template. The detection limit of PCR was assessed by serially diluting known numbers of bacterial cells in sterile distilled water and then subjecting each suspension to PCR. The multiplex PCR mixture contained 0.2 µM of each primer of each group, 10

µl of 2 × Mighty Amp Buffer Ver.2 (Takara Bio Inc., Shiga, Japan), 0.4 µl of Mighty Amp DNA Polymerase (Takara), and 5 µl of the template in a final volume of 20 µl. PCR reactions were performed in a DNA thermal cycler (Applied Biosystems 2720 Thermal Cycler; Applied Biosystems, CA, USA). PCR conditions included an initial denaturation step at 98 °C for 2 min, followed by 30 cycles consisting of 98 °C for 10 s and 70 °C for 1 min. PCR products were analyzed by 2.0% agarose gel electrophoresis before being visualized by electrophoresis in 1 × Tris-borate-EDTA on a 2% agarose gel stained with ethidium bromide. A 100-bp DNA ladder (Takara Biomed, Shiga, Japan) was used as a molecular size marker.


## Results Primer design

Twenty specific primers covering the upstream regions of the 16S rDNA sequences of oral *Actinomyces* species belonging to groups A and B were designed in the present study (Figures 1,2). The specific forward primers of group A were designated as MAJF for *A. johnsonii*, MAORF for *A. oris*, MANF for *A. naeslundii*, MAODF for *A. odontolyticus*, and MAIF for *A. israelii*, whereas


the specific reverse primers were designated as MAJR for *A. johnsonii*, MAORR for *A. oris*, MANR for *A. naeslundii*, MAODR for *A. odontolyticus*, and MAIR for *A. israelii*. The amplicon sizes of *A. johnsonii*, *A. oris*, *A. naeslundii*, *A. odontolyticus*, and *A. israelii* were 136 bp, 294 bp, 413 bp, 622 bp, and 907 bp, respectively. The specific forward primers of group B were designated as AGEOF for *A. georgiae*, ADF for *A. dentalis*, AGERF for *A. gerencseriae*, AGRF for *A. graevenitzii*, and AMF for *A. meyeri*, whereas the specific reverse primers were designated as AGEOR for *A. georgiae*, ADR for *A. dentalis*, AGERR for *A. gerencseriae*, AGRR for *A. graevenitzii*, and AMR for *A. meyeri*. The amplicon sizes of *A. georgiae*, *A. dentalis*, *A. gerencseriae*, *A. graevenitzii*, and *A. meyeri* were 328 bp, 464 bp, 583 bp, 837 bp, and 942 bp, respectively.

## Multiplex PCR

**Detection limit:** Our multiplex PCR method for identifying oral *Actinomyces* species successfully amplified DNA fragments of the expected size for each species (Figures 3,4). The detection limit was assessed in the presence of titrated bacterial cells, and




**Figure 1:** Locations and sequences of species-specific primers for the 16S rDNA of group A oral *Actinomyces* species. The nucleotide sequence of each primer has been underlined.



<p>a: Forward primer for <i>Actinomyces georgiae</i> AGEOF : <u>TCTTCTCCGGGATAACGGTCT</u></p> <p>122 142</p> <p><i>A. georgiae</i> <u>TCTTCTT</u> <u>CGGGATAACGGTCT</u></p> <p><i>A. dentalis</i> <u>CCTGCTC</u> <u>CTGGATAACTGCTT</u></p> <p><i>A. gerencseriae</i> <u>TTACAC</u> <u>CTGGATAACCGCTT</u></p> <p><i>A. graevenitzi</i> <u>CTTACTTTCCTGGATAACTGCTC</u></p> <p><i>A. meyeri</i> <u>TCTTCTT</u> <u>TGGGATAACGCCCC</u></p>		<p>e: Forward primer for <i>Actinomyces gerencseriae</i> AGERF : <u>TGAGGAAGGCCCTTGCCCGTG</u></p> <p>414 433</p> <p><i>A. georgiae</i> <u>GGTCAAGCCGCAACT</u> <u>G</u></p> <p><i>A. dentalis</i> <u>GGTGAAGCCCTGTCCG</u> <u>TTG</u></p> <p><i>A. gerencseriae</i> <u>TGAGGAAGGCCCTTGCCCGT</u> <u>G</u></p> <p><i>A. graevenitzi</i> <u>GAACAAGGCACTGCT</u> <u>TTT</u></p> <p><i>A. meyeri</i> <u>GGTCAAGCCGCAACT</u> <u>G</u></p>		<p>i: Forward primer for <i>Actinomyces meyeri</i> AMF : <u>CTAGCTTGCTGGGTGGATGA</u></p> <p>63 82</p> <p><i>A. georgiae</i> <u>CTGGC</u> <u>TTTGTTGGGT</u> <u>GGATGA</u></p> <p><i>A. dentalis</i> <u>CCGCCCTGTTTTGGGGT</u> <u>GGGTGA</u></p> <p><i>A. gerencseriae</i> <u>CTGGCTTTTTGGTTGGGT</u> <u>TCTGGGGTTA</u></p> <p><i>A. graevenitzi</i> <u>CAAGC</u> <u>TTG</u> <u>CTTGGT</u> <u>GATTA</u></p> <p><i>A. meyeri</i> <u>CTAGC</u> <u>TT</u> <u>GCTGGT</u> <u>GGATGA</u></p>	
<p>b: Reverse primer for <i>Actinomyces georgiae</i> AGEOR : <u>AGTTGCGGCTTGACCACGGG</u></p> <p>449 430</p> <p><i>A. georgiae</i> <u>AGTTGCGGCTTGACCACGGG</u></p> <p><i>A. dentalis</i> <u>GACAGGGCTTACCACCGG</u></p> <p><i>A. gerencseriae</i> <u>GGCAAGGCCTTCTCACTGG</u></p> <p><i>A. graevenitzi</i> <u>AGCAGTGCCTTGTCCCTAG</u></p> <p><i>A. meyeri</i> <u>AGTTGCGGCTTGACCATGAG</u></p>		<p>f: Reverse primer for <i>Actinomyces gerencseriae</i> AGERR : <u>GACCTTCCGCTTCCGAAGGA</u></p> <p>996 977</p> <p><i>A. georgiae</i> <u>CGCGCGCTCTCC</u> <u>GCGC</u></p> <p><i>A. dentalis</i> <u>G</u> <u>CGCCGTCTCC</u> <u>GGCG</u></p> <p><i>A. gerencseriae</i> <u>GACCTTCCGCTTCCGAAGGA</u></p> <p><i>A. graevenitzi</i> <u>TAAACCCTATCTCT</u> <u>AGAG</u></p> <p><i>A. meyeri</i> <u>CACGACGTCTCC</u> <u>GCAC</u></p>		<p>j: Reverse primer for <i>Actinomyces meyeri</i> AMR : <u>ACGCACGTCTCCGCACGCCCA</u></p> <p>1004 984</p> <p><i>A. georgiae</i> <u>GCGCGCTCTCCGCGCCGCCG</u></p> <p><i>A. dentalis</i> <u>CGCCGTCTCC</u> <u>GGCGCCGCC</u></p> <p><i>A. gerencseriae</i> <u>CCCTCCGTCTCCGAAGGACAACC</u></p> <p><i>A. graevenitzi</i> <u>AACCCTATCTCT</u> <u>AGAGCCACG</u></p> <p><i>A. meyeri</i> <u>ACGCACGTCTCCGCACGCCCA</u></p>	
<p>c: Forward primer for <i>Actinomyces dentalis</i> ADF : <u>CCTGCTCCGGATAACTGCTT</u></p> <p>113 133</p> <p><i>A. georgiae</i> <u>TCTTCTT</u> <u>CGGGATAACGGTCT</u></p> <p><i>A. dentalis</i> <u>CCTGCTC</u> <u>CTGGATAACTGCTT</u></p> <p><i>A. gerencseriae</i> <u>TTCACTT</u> <u>CTGGATAACCGCTT</u></p> <p><i>A. graevenitzi</i> <u>CTTACTTTCCTGGATAACTGCTC</u></p> <p><i>A. meyeri</i> <u>TCTTCTT</u> <u>TGGGATAACGCCCC</u></p>		<p>g: Forward primer for <i>Actinomyces graevenitzi</i> AGRF : <u>GACCAGCTGCATGGGTTGG</u></p> <p>146 165</p> <p><i>A. georgiae</i> <u>ACTGGTCTCGCATGG</u> <u>GGG</u></p> <p><i>A. dentalis</i> <u>CGGCTGTCC</u> <u>GCATGGGCTG</u></p> <p><i>A. gerencseriae</i> <u>TGGGTCTGTC</u> <u>GCATGG</u> <u>CGGG</u></p> <p><i>A. graevenitzi</i> <u>G</u> <u>ACCAGCCT</u> <u>GCATGGGTTGG</u></p> <p><i>A. meyeri</i> <u>ACTGGTCTCGCATGG</u> <u>GG</u></p>			
<p>d: Reverse primer for <i>Actinomyces dentalis</i> ADR : <u>GGCCGCTACAAGCCCTTCA</u></p> <p>576 557</p> <p><i>A. georgiae</i> <u>AGCCGCTACAAGCCCTTTA</u></p> <p><i>A. dentalis</i> <u>GGCCGCTACAAGCCCTTCA</u></p> <p><i>A. gerencseriae</i> <u>AACCCGCTACAAGCCCTTTA</u></p> <p><i>A. graevenitzi</i> <u>AGCCGCTACGAGCTTTTA</u></p> <p><i>A. meyeri</i> <u>AACCCGCTACAAGCCCTTTA</u></p>		<p>h: Reverse primer for <i>Actinomyces graevenitzi</i> AGRR : <u>TATCTCTAGAGCCACGCTTAA</u></p> <p>982 962</p> <p><i>A. georgiae</i> <u>CGTCTCC</u> <u>GCGCGCCCGGGGCG</u></p> <p><i>A. dentalis</i> <u>CGTCTCC</u> <u>GGCGCCGCCGCCCC</u></p> <p><i>A. gerencseriae</i> <u>CGTCTCCGAAGGACAACCG</u> <u>GCA</u></p> <p><i>A. graevenitzi</i> <u>TATCTCT</u> <u>AGAGCCACGCTTAA</u></p> <p><i>A. meyeri</i> <u>CGTCTCC</u> <u>GCACGCCACGGGCG</u></p>			


**Figure 2:** Locations and sequences of species-specific primers for the 16S rDNA of group B oral *Actinomyces* species. The nucleotide sequence of each primer has been underlined.



**Figure 3:** Multiplex PCR assay to detect group A oral *Actinomyces* species.

The primer mixture contained MAJF, MAJR, MAORF, MAORR, MANF, MANR, MAODF, MAODR, MAIF, and MAIR.

**Lanes:** 1, *A. johnsonii* JCM 16129; 2, *A. oris* ATCC 27044; 3, *A. naeslundii* ATCC 12104; 4, *A. odontolyticus* ATCC 17929; 5, *A. israelii* ATCC 12102; 6, *A. georgiae* DSM 6843; 7, *A. dentalis* DSM 19115; 8, *A. gerencseriae* JCM 12963; 9, *A. graevenitzi* DSM 15540; 10, *A. meyeri* ATCC 35568; 11, *A. viscosus* ATCC 15987; 12, *Streptococcus oralis* ATCC 10557; 13, *S. salivarius* JCM 5707; 14, *S. anginosus* ATCC 11391; 15, *S. mutans* NCTC 10449; 16, *S. sobrinus* ATCC 33478; 17, *Rothia dentocariosa* JCM 3067; 18, *R. mucilaginoso* JCM 10910; 19, *Corynebacterium matruchotii* ATCC 14266; 20, *C. durum* ATCC 33449; 21, *Neisseria sicca* ATCC 29256; 22, *Aggregatibacter actinomycetemcomitans* ATCC 33384; 23, *Staphylococcus aureus* JCM 28744; 24, *S. epidermidis* ATCC 2414. M, molecular size marker (100-bp DNA ladder).



**Figure 4:** Multiplex PCR assay to detect group B oral *Actinomyces* species.

The primer mixture contained AGEOF, AGEOR, ADF, ADR, AGERF, AGERR, AGRF, AGRR, AMF, and AMR.

**Lanes:** 1, *A. johnsonii* JCM 16129; 2, *A. oris* ATCC 27044; 3, *A. naeslundii* ATCC 12104; 4, *A. odontolyticus* ATCC 17929; 5, *A. israelii* ATCC 12102; 6, *A. georgiae* DSM 6843; 7, *A. dentalis* DSM 19115; 8, *A. gerencseriae* JCM 12963; 9, *A. graevenitzi* DSM 15540; 10, *A. meyeri* ATCC 35568; 11, *A. viscosus* ATCC 15987; 12, *Streptococcus oralis* ATCC 10557; 13, *S. salivarius* JCM 5707; 14, *S. anginosus* ATCC 11391; 15, *S. mutans* NCTC 10449; 16, *S. sobrinus* ATCC 33478; 17, *Rothia dentocariosa* JCM 3067; 18, *R. mucilaginoso* JCM 10910; 19, *Corynebacterium matruchotii* ATCC 14266; 20, *C. durum* ATCC 33449; 21, *Neisseria sicca* ATCC 29256; 22, *Aggregatibacter actinomycetemcomitans* ATCC 33384; 23, *Staphylococcus aureus* JCM 28744; 24, *S. epidermidis* ATCC 2414. M, molecular size marker (100-bp DNA ladder).

the sensitivity of the PCR assay was between  $5 \times 10^3$  and  $5 \times 10^5$  CFU per PCR template (3.6  $\mu$ l) for the *A. johnsonii*-specific primer set with strain JCM 16129, the *A. oris*-specific primer set with strain ATCC 27044, the *A. naeslundii*-specific primer set with strain ATCC 12104, the *A. odontolyticus*-specific primer set with strain ATCC 17929, the *A. israelii*-specific primer set with strain ATCC 12102, the *A. georgiae*-specific primer set with strain DSM 6843, the *A. dentalis*-specific primer set with strain DSM 19115, the *A. gerencseriae*-specific primer set with strain JCM 12963, the *A. graevenitzii*-specific primer set with strain DSM 15540, and the *A. meyeri*-specific primer set with strain ATCC 35568 (data not shown).

**Assay of representative oral bacteria:** As representative oral bacteria, some *Streptococcus*, *Corynebacterium*, *Rothia*, *Neisseria*, *Aggregatibacter*, and *Staphylococcus* were subjected to PCR using the designed primer sets. However, no amplicons were produced from any of the representative oral bacteria (Figures 3,4).

## Discussion and Conclusion

*Actinomyces* is one of the predominant genera in the oral cavity. At the age of 2 months, one-third of infants are already colonized with *Actinomyces* [9]. *Actinomyces* species play a central role in the initial stages of biofilm formation on teeth (i.e., dental plaque) both above (supragingival) and below (subgingival) the gum line [10]. Ten species (i.e., *A. naeslundii*, *A. odontolyticus*, *A. oris*, *A. georgiae*, *A. gerencseriae*, *A. graevenitzii*, *A. dentalis*, *A. johnsonii*, *A. israelii*, and *A. meyeri*) among the genus *Actinomyces* are regarded as normal human oral bacteria [3].

*Actinomyces* species are mainly associated with cervicofacial actinomycosis, oral or cerebral abscesses, caries, root canal infections and periodontitis [11-13]. These microorganisms appear to play a more important role than expected in the pathogenesis of osteoradionecrosis- and bisphosphonate-related osteonecrosis of the jaw [14-16], and may cause lethal infections such as mediastinitis [17]. As a consequence, fast and reliable identification methods for these microorganisms have become increasingly important.

The isolation and identification of *Actinomyces* using conventional methods are often difficult and time-consuming. Previous studies were conducted in order to characterize them using phenotypic [18-20] and molecular [21,22] approaches. Most of the available commercial identification kits do not encompass the majority of new species in their database.

The PCR method is a molecular technique that allows species verification at the genetic level, and possesses high sensitivity and bacterial species specificity. It is also useful for the detection and identification of fungi [23]. Moreover, it may efficiently and accurately detect specific microorganisms in clinical samples, regardless of whether they are alive or dead. Also, it might be an innovative method for detection of microorganisms difficult to culture using traditional microbiological techniques. The

16S rDNA region of the bacterial genome provides an ideal target for species identification using PCR [16]. Moreover, multiplex-PCR is a rapid tool that allows for the simultaneous amplification of more than one sequence of target DNA in a single reaction, thereby saving time and reagents [24]. On the other hand, Hirotsuki et al. [25] and we [26,27] developed the multiplex-PCR, which allows for the amplification of one sequence of target DNA in a single reaction containing the plural species-specific primers, for bacterial species identification. Previous studies have used different molecular methodologies to identify and differentiate *Actinomyces* from oral samples after anaerobic cultivation, including PCR-RFLP, chromosomal DNA fingerprinting, 16S rRNA gene sequencing and oligonucleotide-DNA hybridization using universal primers or oligonucleotide probes [13,28-30]. Tian et al. reported rapid and sensitive PCR-dipstick DNA chromatography assay that could be read by eye for multiplex and semi quantitative analysis of dental plaque bacteria including *Actinomyces* species [31].

*In silico* method helps in designing primers, and there are various programs available for PCR primer design [32]. In the present study, we designed species-specific primers with the already mentioned means, for the identification of oral *Actinomyces* species using a PCR method. These primers were able to distinguish each oral *Actinomyces* species and did not display cross-reactivity with representative oral bacteria or other *Actinomyces* species. Moreover, we developed a multiplex PCR method with the ability to identify and differentiate oral *Actinomyces* species (i.e., *A. naeslundii*, *A. johnsonii*, *A. oris*, *A. odontolyticus*, *A. israelii*, *A. georgiae*, *A. dentalis*, *A. graevenitzii*, *A. gerencseriae*, and *A. meyeri*) using only two PCR tubes per sample.

Our multiplex PCR method is easy because the use of Mighty Amp DNA Polymerase Ver.2 (Takara) means that DNA extraction may be avoided, and species identification using this method only takes approximately 2 hours. Thus, the method described herein will allow the prevalence of oral *Actinomyces* species and their involvement in oral infections to be fully clarified in future studies.

## References

1. Sabbe LJ, Van de Merwe D, Schouls L, Bergmans, A, Vaneechoutte M, et al. (1999) Clinical spectrum of infections due to the newly described *Actinomyces* species *A. turicensis*, *A. radingae*, and *A. europaeus*. *J Clin Microbiol* 37: 8-13.
2. Schaal KP, Lee HJ (1992) Actinomycete infections in humans-a review. *Gene* 115: 201-211.
3. Könönen E, Wade WG (2015) *Actinomyces* and related organisms in human infections. *Clin Microbiol Rev* 28: 419-442.
4. Socransky SS, Hubersak C, Propas D (1970) Induction of periodontal destruction in gnotobiotic rats by a human oral strain of *Actinomyces naeslundii*. *Arch Oral Biol* 15: 993-995.
5. Dzik JL, Socransky SS, Haffajee AD (1988) The predominant cultivable microbiota of active and inactive lesions of destructive periodontal diseases. *J Clin Periodontol* 15: 316-323.

6. Liljemark WF, Bloomquist CG, Bandt CL, Pihlstrom BL, Hinrichs JE, et al. (1993) Comparison of the distribution of *Actinomyces* in dental plaque on inserted enamel and natural tooth surfaces in periodontal health and disease. *Oral Microbiol Immunol* 8: 5-15.
7. Moore LV, Moore WE, Cato EP, Smibert RM, Burmeister JA, et al. (1987) Bacteriology of human gingivitis. *J Dent Res* 66: 989-995.
8. Putnins EE, Bowden GH (1993) Antigenic relationships among oral *Actinomyces* isolates *Actinomyces naeslundii* genospecies 1 and 2, *Actinomyces howellii*, *Actinomyces denticolens*, and *Actinomyces slackii*. *J Dent Res* 72: 1374-1385.
9. Sarkonen N, Könönen E, Summanen P, Kanervo A, Takala A, et al. (2000) Oral colonization with *Actinomyces* species in infants by two years of age. *J Dent Res* 79: 864-867.
10. Zijngje V, van Leeuwen MBM, Degener JE, Abbas F, Thurnheer T, et al. (2010) Oral biofilm architecture on natural teeth. *PLoS One* 5: e9321.
11. Marsh P, Martin MV (1999) *Oral microbiology*, fourth edn. Oxford: Reed Educational and Professional Publishing.
12. Brailsford SR, Tregaskis RB, Leftwich HS, Beighton D (1999) The predominant *Actinomyces* spp. isolated from infected dentin of active root canal lesions. *J Dent Res* 78: 1525-1534.
13. Tang G, Samaranyake LP, Yip HK, Chu FC, Tsang PC, et al. (2003) Direct detection of *Actinomyces* spp. from infected root canals in a Chinese population: a study using PCR-based, oligonucleotide-DNA hybridization technique. *J Dent* 31: 559-568.
14. Hansen T, Kunkel M, Kirkpatrick CJ, Weber A (2006) *Actinomyces* in infected osteoradionecrosis - underestimated? *Hum Pathol* 37: 61-67.
15. De Ceulaer J, Tacconelli E, Vandecasteele SJ (2014) *Actinomyces* osteomyelitis in bisphosphonate-related osteonecrosis of the jaw (BRONJ): the missing link? *Eur J Clin Microbiol Infect Dis* 33: 1873-1880.
16. Panya S, Fliefel R, Probst F, Tröltzsch M, Ehrenfeld M, et al. (2017) Role of microbiological culture and polymerase chain reaction (PCR) of *actinomyces* in medication-related osteonecrosis of the jaw (MRONJ). *J Craniomaxillofac Surg* 45: 357-363.
17. Branquinho DF, Andrade DR, Almeida N, Sofia C (2014) Mediastinitis by *Actinomyces meyeri* after oesophageal stent placement. *BMJ Case Rep*.
18. Brander MA, Jousimies-Somer HR (1992) Evaluation of the RapID ANA II and API ZYM systems for identification of *Actinomyces* species from clinical specimens. *J Clin Microbiol* 30: 3112-3116.
19. Sarkonen N, Könönen E, Summanen P, Könönen M, Jousimies-Somer H (2001) Phenotypic identification of *Actinomyces* and related species isolated from human sources. *J Clin Microbiol* 39: 3955-3961.
20. Kerttula AM, Carlson P, Sarkonen N, Hall V, Könönen E (2005) Enzymatic/biochemical analysis of *Actinomyces* with commercial test kits with an emphasis on newly described species. *Anaerobe* 11: 99-108.
21. Hall V (2008) *Actinomyces* - gathering evidence of human colonization and infection. *Anaerobe* 14: 1-7.
22. Henssge U, Do T, Radford DR, Gilbert SC, Clark D, et al. (2009) Emended description of *Actinomyces naeslundii* and descriptions of *Actinomyces oris* sp. nov. and *Actinomyces johnsonii* sp. nov., previously identified as *Actinomyces naeslundii* genospecies 1, 2 and WVA 963. *Int J Syst Evol Microbiol* 59: 509-516.
23. Al-Shuhaib MBS, Albakri AH, Alwan SH, Almandil NB, AbdulAzeez S, et al. (2018) Optimal pcr primers for rapid and accurate detection of *Aspergillus flavus* isolates. *Microbial pathog* 116: 351-355.
24. Henegariu O, Heerema NA, Dlouhy SR, Vance GH, Vogt PH (1997) Multiplex PCR: critical parameters and step-by-step protocol. *Biotechniques* 23: 504-511.
25. Hirotaki S, Sasaki T, Kuwahara-Arai K, Hiramatsu K (2011) Rapid and accurate identification of human-associated staphylococci by use of multiplex PCR. *J Clin Microbiol* 49: 3627-3631.
26. Tsuzukibashi O, Uchibori S, Shinozaki-Kuwahara N, Kobayashi T, Takada K, et al. (2014) A selective medium for the isolation of *Corynebacterium* species in oral cavities. *J Microbiol Methods* 104: 67-71.
27. Tsuzukibashi O, Uchibori S, Kobayashi T, Umezawa K, Mashimo C, et al. (2017) Isolation and identification methods of *Rothia* species in oral cavities. *J Microbiol Methods* 134: 21-26.
28. Sato T, Matsuyama J, Takahashi N, Sato M, Johnson J, et al. (1998) Differentiation of oral *actinomyces* species by 16s ribosomal DNA polymerase chain reaction-restriction fragment length polymorphism. *Arch Oral Biol* 43: 247-252.
29. Ruby JD, Li Y, Luo Y, Caufield PW (2002) Genetic characterization of the oral *actinomyces*. *Arch Oral Biol* 47: 457-463.
30. Tang G, Samaranyake LP, Yip HK (2004) Genotypic diversity of oral *Actinomyces naeslundii* genospecies 1 and 2 in caries-active preschool children. *Oral Microbiol Immunol* 19: 371-378.
31. Tian L, Sato T, Niwa K, Kawase M, Tanner AC, et al. (2014) Rapid and Sensitive PCR-Dipstick DNA Chromatography for Multiplex Analysis of the Oral Microbiota. *BioMed Research International*.
32. Kumar A, Chordia N (2015) In silico PCR primer designing and validation. *Methods Mol Biol* 1275: 143-151.