

Regulation Studies of a Cuticle Protein Underlying Genomic Analysis

Md. Saheb Ali^{1†}, Birendra Mishra², Mohammad Sahin Polan^{1†}, Osamu Ninagi¹, Ahsanul Haque Swapon³ and Masamitsu Yamaguchi⁴

¹Agriculture Wing, Bangladesh Jute Research Institute, Manik Mia Ave, Dhaka, Bangladesh

²Department of Medicine, University of California -Irvine, Irvine, CA 92697, USA

³Entomology Department, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh

⁴Department of Applied Biology, Kyoto Institute of Technology, Sakyo-ku, Kyoto 606-8585, Japan

[†]United Graduate School of Agricultural Science, Tokyo University of Agriculture and Technology, Tokyo, Japan

Corresponding author: Md. Saheb Ali, Agriculture Wing, Bangladesh Jute Research Institute, Manik Mia Ave, Dhaka 1207, Bangladesh, Tel: +88-01714538138; **E-mail:** sahebbjri@yahoo.com

Received date: 16 Dec 2015; **Accepted date:** 05 Jan 2016; **Published date:** 10 Jan 2016.

Citation: Ali MS, Mishra B, Polan MS, Ninagi O, Swapon AH, et al. (2016) Regulation Studies of a Cuticle Protein Underlying Genomic Analysis. *Int J Mol Genet Gene Ther* 2(1): doi <http://dx.doi.org/10.16966/2471-4968.104>

Copyright: © 2016 Ali MS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The present study was aimed to understand the regulatory mechanism of a cuticular protein gene coding for cuticle protein, *CPR95* of *Bombyx mori* metamorphosis. Developmental profile of *CPR95* resembled that of *E74A*. *E74A* and *CPR95* transcripts induced by the 20E pulse treatment *in vitro*. Therefore, the results showed the relatedness of *CPR95* and *E74A*. Site-directed mutagenesis of *E74A* binding site in the upstream of *CPR95* and a reporter assay demonstrated the strong relatedness of *CPR95* and *E74A*. Through genomic information and a transient reporter assay, we proved the regulating mechanism of *CPR95* expression. The present findings showed that the temporal expression of *CPR95* was regulated by ecdysone-responsive transcription factor *E74A* at the pre-pupal stage in the wing discs of *Bombyx mori*.

Keywords: Temporal expression; Cuticular protein; Ecdysone pulse; Site-directed mutagenesis; Gene gun

Introduction

Insect cuticle is a multi-layered structure with three functional regions: epicuticle, procuticle, and endocuticle, which differs in protein composition, structural features and physiological functions [1]. The character of insect cuticle is determined by the construction of cuticular protein, and the various cuticle types show pronounced differences in mechanical properties and these differences are related to the properties of the individual proteins [2]. The character of insect cuticle differs in species and stages, and the difference is brought about by the combination of cuticular proteins. Cuticular proteins are characterized by repeated occurrence of a few small motifs consisting mainly of hydrophobic residues [2]. Thus, insects make up new cuticle at the each larval molting and different types of cuticle at the larval to pupal or pupal to adult transformation. Insect cuticle is mainly made up of chitin and cuticle proteins, and cuticular proteins determine the character of cuticle and bring about the variety of insect cuticle.

Ecdysone-responsive expression of cuticular proteins has been reported [3-12]. Most of their expressions were induced by an ecdysteroid pulse; the expression required the existence and removal of 20E [9,11,13]. This condition is similar to that of the stage around the ecdysis. Few examples were reported to be up-regulated by 20E [7,9,10]. They were induced by the addition of ecdysone *in vitro*. Thus, cuticular protein genes, which have different developmental profiles, were induced by different types of ecdysone signal, and these things were described in a recent review [14].

Ecdysone-responsive transcription factors have been shown to determine the expression of cuticular protein genes [4]. β FTZ-F1 increased the promoter activity of the cuticular protein gene, *BMWCP5*, which was expressed around pupation when the ecdysteroid titer decreased after its peak [15,16]. BR-C Z2 activated the *BmorCPG11* promoter independently from other ecdysone responsive transcription factors [6]. In addition to above transcription factors, EcR bound to the promoter region of the cuticular protein gene, *BMWCP10*, and activated its transcription [15]. BHR3 regulated *BmorCPH5*, and their transcription

peak was observed earlier than that of *BMWCP5* [4]. Thus, transcription factors bind to upstream of the target cuticular protein genes and bring about the stage-specific expression of cuticular protein genes in wing discs and epidermis. Therefore, cuticular protein genes are targets of ecdysone-responsive transcription factors, as described above. These studies have been conducted by using a genomic database and hypothesized that the ecdysone-responsive transcription factor, *E74A*, regulates the expression of *CPR95*.

Materials and Methods

Experimental animals

A hybrid strain of *B. mori* was reared at 25°C in a 12 h light: 12 h dark photoperiod. Larvae began wandering on six day of the fifth larval instar, pupation occurred 3 days thereafter, and adults enclosed 10 days after pupation. The periods (in days) corresponding to the developmental stages of the fourth to fifth larval ecdysis, wandering, pupation, and eclosion were designated as V0, W0, P0, and A0, respectively. The three days before pupation were designed as W1-W3. The W3 stage was divided into three different sub-stages, W3 early (W3E), W3 mid (W3M), and W3 late (W3L). The W3 sub-stages were determined on the time and visible shortening of the length of the leg [3].

In vitro culture of wing discs

Wing discs of larvae at W2 stages were prepared and cultured *in vitro* as previously described [4]. The culture was carried out at 25°C under sterile conditions according to a previous report [17].

Quantitative reverse transcription-PCR

To determine expression levels of cuticular protein gene and transcription factor, total RNA and cDNA synthesis were done according to a previous report [4]. Quantitative reverse transcription-PCR was performed using an ABI7500 real-time PCR machine (Applied Biosystems) using Fast Start Universal SYBR Green Master (Roche) according to the manufacturer's protocol. The data were normalized in

each sample using gene for the ribosomal protein S4. The oligonucleotide primer sets were designed using Primer3 software (<http://frodo.wi.mit.edu/>) as listed in Table 1.

Plasmid construction and mutagenesis

Upstream regions (-2040 to +18 and -136 to +18) of *CPR95* were amplified by PCR according to a previous report [3]. The amplified DNA fragments were digested with *SacI* and *NheI* and then ligated into the *SacI* and *NheI* sites of the luciferase reporter plasmid pGL3-basic to generate the constructs. A *Renilla* luciferase reporter (*PhRG-hsp*) driven by the *Drosophila* heat-shock protein70 promoter [18] was used as a normalization control [11]. Single mutant, (-104/-102) E74A mut for *CPR95* was performed with the Quick-Change™ Site-Directed Mutagenesis Kit (Stratagene) using the *CPR95-136* plasmid as the template. Twenty nanograms of plasmid DNA were amplified with *Pfu* DNA polymerase followed by digestion of the parental plasmid by *DpnI*. The mutated plasmid was introduced into XL1-Blue super-competent cells. The mutagenic reactions were performed according to the manufacturer's instructions. The introduction of each mutation was confirmed by sequencing. Oligonucleotide primers used to generate the constructs and mutation are shown in Table 1.

Luciferase transient expression assays

Transient expression of the reporter constructs in the wing discs was performed as previously described [4]. Following manufacturer's instructions, 12.5 mg of gold particles (diameter: 1.0 μm) was coated with plasmid DNA (50 μg each of pGL3-derived vectors and 5 μg *phRG-hsp*). The reporter constructs were introduced into wing discs using a particle gun (Bio-rad). Bombardment proceeded under helium pressure of 150 psi (pounds per square inch). After bombardment, wing discs were cultured for 48 h at 25°C in Grace's medium (Invitrogen) with or without 2 μg/ml of 20E (Sigma). The culture method was described previously [17]. After 48 h of culture, wing discs were washed twice in PBS. The tissues were suspended in a 25 μl 1x reporter Passive Lysis Buffer (Promega) and frozen/thawed for 5 cycles in liquid nitrogen before the supernatant was equilibrated at 4°C for 1 h. The supernatant was collected by centrifugation at 12,000 g for 2 min at 4°C. The luciferase reporter assay was carried out using a Dual-Luciferase reporter assay system (Promega) in a luminometer (Perkin Elmer) according to the manufacturer's protocol. The luciferase activity was normalized to the level of the *Renilla* luciferase activity. All experiments were performed at least five times. The results were expressed as the mean ± S. E. M., and significance was set at $p < 0.05$.

Histological study

Pieces of integument containing the head, thorax and abdomen (Figure 1) were dissected, fixed in Carnoy's fixative for 2 h, and then dehydrated and embedded in paraffin. Seven-micrometer sections were de-waxed in

xylene, rehydrated through an alcohol series, and then washed with tap water for 15 min. After washing, a Periodic acid Schiff (PAS) reaction [19] was performed to detect the polysaccharide derived from BM. The slides were then dipped in Lillie solution for 10 min (100 ml KIO_4 + 0.4 ml of 60% HNO_3), in Schiff's reagents for 30 min, and then in a H_2SO_3 solution [5 ml of 1N HCl + 5 ml of 10% $K_2S_2O_5$ + 100 ml H_2O] (3×1 min), and rinsed with tap water for 10 min. The sections were then stained with hematoxylin for 5 min, destained via immersion in acid alcohol (70% ethanol + 1 ml 1N HCl) for 1 min, and washed in running tap water for 15 min. After washing, they were dehydrated in a graded series of ethanol (70%, 90%, 95%, and absolute), cleared in xylene, and mounted using Canada balsam and a cover slip. The sections were visualized under a light microscope (Olympus BX50) and documented using DP-BSW software.

Results

CPR95 showed distinct expression profile among the cuticular protein genes expressed in wing discs of *Bombyx mori*

A cuticular protein gene, *CPR95* (Figure 2A), expressed in the fifth larval stage of *Bombyx mori*, which was similar to that of *E74A*. *E74A* is one of ecdysone-responsive transcription factor and showed expression peak at the W3L stage when ecdysteroid in the hemolymph declined at the pre-pupal period. Therefore, to understand the expression profile of *E74A* and *CPR95*, we analyzed the expression of *E74A* and *CPR95*.




Figure 1: *Bombyx* larvae of W3 mid stage. A. An arrow indicates the dissected position for samples of thoracic epidermis. B. Arrows indicate the positions for samples of abdominal epidermis (A), wing disc (W) and head capsule (H).

Gene name	Accession Number	Forward Primer	Reverse Primer	Size of Amplicon (bp)
<i>CPR95</i>	BR000596	5'- AGTTTCCTCCAGAGCATCG-3'	5'-AGACGAATAGTGGACTGGAGC-3'	145
<i>E74A</i>	DQ471939	5'- GCACCACCTATCGAGATAAAGC-3'	5'- CTGCCCGTTGTTTGTAAATG-3'	117
<i>rpl</i>	NM_001043792	5'-GATTCACAATCCACCGTATCACC-3'	5'-CCATCATGCGTTACCAAGTACG-3'	109
The oligonucleotide primers used to generate different constructs were as follows (The restriction enzymes sites are underlined)				
<i>CPR95-2040</i>		5'- <u>CCCGAGCTC</u> CGCGGACAGATACCACCGCCCTGCC-3'		
<i>CPR95-136</i>		5'- <u>CCCGAGCTC</u> GAATGGATTGATTCACGAAACTATG--3'		
<i>CPR95-+18</i>		5'- <u>CCCGCTAGC</u> GATCAACTGTTGAATGATGCTTATG-3'		
The oligonucleotide primer used to generate the mutation was shown as follow (The underlined bases were mutated)				
<i>CPR95</i> (-104/-102) E74A mut		5'-CTATGTGACAAG <u>T</u> AGCAAGGTCACCATAAGCTGTC-3'		

Table 1: List of Primers




Figure 2: Developmental profile of cuticular protein gene *CPR95* (A) and ecdysone responsive transcription factor *E74A* (B). RNA was extracted from wing discs and reverse-transcribed to cDNA for use in Real-Time PCR. Values represent the mean \pm S.E.M. of results from three independent experiments.




Figure 3: Effect of ecdysone pulse treatment on *E74A* (A) and *CPR95* (B). RNA was extracted from wing discs and reverse-transcribed to cDNA for use in Real-Time PCR. Wing discs of the W2 stage were incubated 12 h in a medium containing 2 μ g/ml 20E and then transferred to a hormone-free medium with or without cycloheximide (50 μ g/ml) for the indicated time. Values represent the mean \pm S.E.M. of results from three independent experiments. Asterisks indicate $p < 0.05$ significance by the student's t-test.

The expression profile of *CPR95* and *E74A* is similar in wing discs

The expression of *CPR95* increased rapidly after W3E stage and peaked at W3L stage, when the ecdysteroid titer in the hemolymph declining, then it rapidly decreased (Figure 2A). *E74A* transcripts increased rapidly after W3E stage and peaked at W3L stage then it rapidly decreased (Figure 2B), which is similar to the expression of *CPR95*. *E74A* and *CPR95* transcripts rapidly decreased after W3L stage (Figure 2), when the hemolymph ecdysteroid titer decreased. Therefore the effect of ecdysone removal after treatment (ecdysone pulse) was examined. Transcripts of *E74A* increased after hormone removal (Figure 3A), which were not observed by the addition of cycloheximide. Transcripts of *CPR95* also showed a similar increase after 20E removal (Figure 3B), and the addition of cycloheximide inhibited the transcription. The results suggest that the transcription of both genes was activated with the 20E removal. Both genes showed similar ecdysone pulse responsiveness; transcription was induced by the removal of 20E. From the similarity of expression profiles, it is suggested

that *CPR95* is induced by the ecdysone responsive transcription factor, *E74A*. Therefore we searched the upstream genomic sequences of *CPR95* and found two putative *E74A* binding sites [4] upstream of *CPR95* gene (Figure 4). Then, we applied reporter assay for the promoter analysis of *CPR95* related with *E74A*.

Reporter assay showed the relatedness of *CPR95* and *E74A*

A plasmid containing a 2040 bp region upstream of *CPR95* showed a higher promoter activity by the 20E pulse treatment (Figure 4). A deleted construct (*CPR95-136*) containing a putative *E74A* binding site showed a similar activity (Figure 4). Therefore, we conducted mutagenesis on this binding site using *CPR95-136*, resulted in the reduction of promoter activity (Figure 4). Luciferase activity of *CPR95* promoter region in the media without 20E decreased to that cultured with 20E. This suggests mutation abolished the effect of 20E pulse that is suggested to be the effect of *E74A*. This result strongly suggests that the promoter activity of *CPR95* was regulated by *E74A*.

Discussion

Different expression patterns of cuticular protein genes in the wing discs of the fifth larval instar have reported and clarified the regulation of cuticular protein genes of *Bombyx mori* through the genomic database [4,11,15,20]. Here, we have clarified the related ecdysone-responsive transcription factors. The promoter region of the different cuticular protein genes was bound and activated by EcR and different ecdysone-responsive transcription factors, β FTZ-F1, E74A, BR-C Z2, and BR-C Z4, resulting in their different expression patterns. These studies succeeded by using genomic information and a transient reporter assay by using wing discs.

In the present study, we found a cuticular protein gene, *CPR95*, which transcripts increased rapidly after the W3E stage and peaked at W3L, then rapidly decreased. Most cuticular protein genes showed expression peak at around the pupation [21], and transcription factors, β FTZ-F1, BR-C and E74A regulated the expression of these cuticular protein genes [4-6,15]. E74A was the only transcription factor that showed similar developmental profile to *CPR95*. *CPR95* was induced by ecdysone pulse, and showed

similar ecdysone responsiveness to E74A (Figure 3). E74A was induced by ecdysone pulse as previously reported in *D. melanogaster* [22], *M. sexta* [23] and *Bombyx mori* [4,24]. In the present study, we found the relatedness of *CPR95* and E74A. *CPR95* promoter was strongly regulated by the ecdysone pulse treatment, which disappeared by the mutagenesis of proximal putative E74A binding site (Figure 4). The result suggests the strong relatedness of E74A with *CPR95*. Together with the result of qRT-PCR, we concluded that *CPR95* transcription is regulated by E74A. It is suggested that the responsiveness to ecdysone determines the expression timing of transcription factors and their related cuticular protein genes, resulted in the difference of *CPR95* and *BmorCPH5* expression [4]. Cuticular proteins containing R&R residue [25] are reported to bind with chitin [26,27] and would construct the procuticle. Although some of non R&R cuticular-proteins have the cuticle-binding capacity [28,29], most of them have been reported not to bind with chitin [30,31]. Most cuticular protein genes expressed in the wing discs are transcribed before pupation by the signal of ecdysone pulse, and most of them have R&R residue. *CPR95* would construct procuticle, since it has R&R residue and is transcribed later than *BmorCPH5* [4].





Figure 4: Promoter activities of different lengths of the *CPR95* 5'-flanking region. Luciferase reporter constructs having various lengths of the 5'-flanking of *CPR95* wild and mutated constructs were bombarded into wing discs of the W2 stage and cultured for 48 h with or without 2 μ g/ml 20E. Numbers refer to the nucleotide position of the putative transcription start site. The luciferase activity was normalized to the Renilla luciferase activities. Luciferase assays were performed in five times, and results are reported as mean \pm S.E.M. Firefly/Renilla ratios.



The expression of *BR-Z2*, *βFTZ-F1* and *E-74A* was distinct in the epidermis of the head, thorax and abdomen respectively [5,6]. Histological photograph and the gene expression pattern demonstrated that different ecdysone-responsive transcription factors were expressed in the different region of the epidermis, which determined the cuticle protein genes expressed there (Figure 5). Thus, these ecdysone-responsive transcription factors are suggested to regulate their target genes, and the series of their expression would bring about insect metamorphosis.

The present findings suggest that cuticular protein genes are expressed in series according to their regulating transcription factors, resulting in a continuous series of cuticular protein production, which enable to construct the epi-, exo-, and endo-cuticle. These different types of cuticle proteins are combined and form the pupal cuticle, and the present findings suggest that ecdysone responsive transcription factors determine the space where cuticular protein genes are expressed. Genomic information and a transient reporter assay will further clarify the mechanism of insect cuticle construction.

References

- Willis JH (1996) Metamorphosis of the cuticle, its proteins, and their genes. In: Gilbert LI, Tana JR, Atkinson BG (Eds.), *Cell Biology: a Series of Monographs: Metamorphosis. Postembryonic Reprogramming of Gene Expression in Amphibian and Insect Cells*, Academic Press, San Diego, 253–282.
- Andersen SO, Højrup P, Roepstorff P (1995) Insect cuticular proteins. *Insect Biochem Mol Bio* 25: 153-176.
- Ali MS, Rahman RF, Swapon AH (2015) Transcriptional Regulation of Cuticular Protein Glycine-Rich13 Gene Expression in Wing Disc of *Bombyx mori*, Lepidoptera. *J Insect Sci* 27: 1-6.
- Ali MS, Iwanaga M, Kawasaki H (2013) Ecdysone-responsive transcriptional regulation determines the temporal expression of cuticular protein genes in wing discs of *Bombyx mori*. *Gene* 512: 337-347.
- Ali MS, Iwanaga M, Kawasaki H (2012) Ecdysone-responsive transcription factors determine the expression region of target cuticular protein genes in the epidermis of *Bombyx mori*. *Dev Genes Evol* 222: 89-97.
- Ali MS, Wang HB, Iwanaga M, Kawasaki H (2012) Expression of cuticular protein genes, *BmorCPG11* and *BMWCP5* is differently regulated at the pre-pupal stage in wing discs of *Bombyx mori*. *Comp Biochem Physiol B Biochem Mol Biol* 162: 44-50.
- Horodyski FM, Riddiford LM (1989) Expression and hormonal control of a new larval cuticular multigene family at the onset of metamorphosis of the tobacco hornworm. *Dev Biol* 132: 292-303.
- Hiruma K, Hardie J, Riddiford LM (1991) Hormonal regulation of epidermal metamorphosis in vitro: control of expression of a larval-specific cuticle gene. *Dev Biol* 114: 369-378.
- Braquart C, Bouhin H, Quenedey A, Delachambre J (1996) Up-regulation of an adult cuticular gene by 20-hydroxyecdysone in insect metamorphosing epidermis cultured in vitro. *Eur J Biochem* 240: 336-341.
- Noji T, Ote M, Takeda M, Mita K, Shimada T, et al. (2003) Isolation and comparison of different ecdysone-responsive cuticle protein genes in wing discs of *Bombyx mori*. *Insect Biochem Mol Biol* 33: 671-679.
- Nita M, Wang HB, Zhong YS, Mita K, Iwanaga M, et al. (2009) Analysis of ecdysone-pulse responsive region of *BMWCP2* in wing disc of *Bombyx mori*. *Comp Biochem Physiol B Biochem Mol Biol* 153: 101-108.
- Wang HB, Iwanaga M, Kawasaki H (2009) Activation of *BMWCP10* promoter and regulation by *BR-C Z2* in wing disc of *Bombyx mori*. *Insect Biochem Mol Biol* 39: 615-623.
- Zhong YS, Mita K, Shimada T, Kawasaki H (2006) Glycine-rich protein genes, which encode a major component of the cuticle, have different developmental profiles from other cuticle protein genes in *Bombyx mori*. *Insect Biochem Mol Biol* 36: 99-110.
- Charles JP (2010) The regulation of expression of insect cuticle protein genes. *Insect Biochem Mol Biol* 40: 205-213.
- Wang HB, Nita M, Iwanaga M, Kawasaki H (2009) *βFTZ-F1* and *Broad-Complex* positively regulate the transcription of the wing cuticle protein gene, *BMWCP5*, in wing discs of *Bombyx mori*. *Insect Biochem Mol Biol* 39: 624-633.
- Sakurai S, Kaya M, Satake S (1998) Hemolymph ecdysteroid titer and ecdysteroid-dependent development events in the last-larval stadium of the silkworm, *Bombyx mori*: role of low ecdysteroid titer in larval-pupal metamorphosis and a reappraisal of the head critical period. *J Insect Physiol* 44 867-881.
- Kawasaki H (1989) Methods for culture of *Bombyx mori* wing discs. *J Tiss Culture Methods* 12: 31-33.
- Tomita S, Kawai Y, Woo SD, Kamimura M, Iwabuchi K, et al. (2001) Ecdysone-inducible foreign gene expression in stably-transformed lepidopteran insect cells. In *Vitro Cell Dev Biol Anim* 37: 564-571.
- Kanayama H, Togami M, Adachi N, Fukai Y, Okumoto T (1986) [Studies on the antitumor active polysaccharides from the mycelia of *Poria cocos* Wolf. III. Antitumor activity against mouse tumors]. *Yakugaku Zasshi* 106: 307-312.
- Wang HB, Moriyama M, Iwanaga M, Kawasaki H (2010) Ecdysone directly and indirectly regulates a cuticle protein gene, *BMWCP10*, in the wing disc of *Bombyx mori*. *Insect Biochem Mol Biol* 40: 453-459.
- Futahashi R, Okamoto S, Kawasaki H, Zhong YS, Iwanaga M, et al. (2008) Genome-wide identification of cuticular protein genes in the silkworm, *Bombyx mori*. *Insect Biochem Mol Biol* 38: 1138-1146.
- Karim DF, Thummel SC (1991) Ecdysone coordinates the timing and amounts of *E74A* and *E74B* transcription in *Drosophila*. *Genes Dev* 5: 1067-1079.
- Stilwell GE, Nelson CA, Weller J, Cui H, Hiruma K (2003) *E74* exhibits stage-specific hormonal regulation in the epidermis of the tobacco hornworm, *Manduca sexta*. *Dev Biol* 258: 76-90.
- Sekimoto T, Iwami M, Sakurai S (2007) 20-hydroxyecdysone regulation of two isoforms of the Ets transcription factor *E74* gene in programmed cell death in the silkworm anterior silk gland. *Insect Mol Biol* 16: 581-590.
- Rebers JE, Riddiford LM (1988) Structure and expression of a *Manduca sexta* larval cuticle gene homologous to *Drosophila* cuticle genes. *J Mol Biol* 203: 411-423.
- Rebers JE, Willis JH (2001) A conserved domain in arthropod cuticular proteins bind chitin. *Insect Biochem Mol Biol* 31: 1083-1094.
- Togawa T, Nakato H, Izumi S (2004) Analysis of the chitin recognition mechanism of cuticle proteins from the soft cuticle of the silkworm, *Bombyx mori*. *Insect Biochem Mol Biol* 34: 1059-1067.
- Tang L, Liang J, Zhan Z, Xiang Z, He N (2010) Identification of the chitin-binding proteins from the larval proteins of silkworm, *Bombyx mori*. *Insect Biochem Mol Biol* 40: 1077-1087.
- Willis H (2010) Structural cuticular proteins from arthropods: annotation, nomenclature, and sequence characteristics in genomics era. *Insect Biochem Mol Biol* 40: 241-251.
- Andersen SO (2000) Studies on proteins in post-ecdysial nymphal cuticle of locust, *Locusta migratoria*, and cockroach, *Blaberus craniifer*. *Insect Biochem Mol Biol* 30: 569-577.
- Togawa T, Augustine Dunn W, Emmons AC, Willis JH (2007) *CPF* and *CPFL*, two related gene families encoding cuticular proteins of *Anopheles gambiae* and other insects. *Insect Biochem Mol Biol* 37: 675-688.